



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C12N 15/55, C07K 16/40,</b> <b>C12N 9/12, G01N 33/50</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/08180</b> <b>(43) International Publication Date:</b> 17 February 2000 (17.02.2000)
<b>(21) International Application Number:</b> PCT/US99/17630 <b>(22) International Filing Date:</b> 03 August 1999 (03.08.1999) <b>(30) Priority Data:</b> 60/095,270 04 August 1998 (04.08.1998) US 60/099,972 11 September 1998 (11.09.1998) US <b>(60) Parent Application or Grant</b> IMMUNEX CORPORATION [/]; (). VIRCA, G., Duke [/]; (). BIRD, Timothy, A. [/]; (). ANDERSON, Dirk, M. [/]; (). MARKEN, John, S. [/]; (). VIRCA, G., Duke [/]; (). BIRD, Timothy, A. [/]; (). ANDERSON, Dirk, M. [/]; (). MARKEN, John, S. [/]; (). JONES, Simone, L. ; ().	<b>Published</b> <i>#9 attached 09/98 474</i>	
<b>(54) Title: HUMAN cDNAs ENCODING POLYPEPTIDES HAVING KINASE FUNCTIONS</b> <b>(54) Titre: ADNc HUMAINS CODANT DES POLYPEPTIDES A FONCTIONS KINASES</b>  <b>(57) Abstract</b> <p>The invention is directed to purified and isolated human polypeptides having kinase function, the nucleic acids encoding such polypeptides, processes for production of recombinant forms of such polypeptides, antibodies generated against these polypeptides, fragmented peptides derived from these polypeptides, the use of such polypeptides and fragmented peptides in phosphorylation reactions and as molecular weight markers, the use of such polypeptides and fragmented peptides as controls for peptide fragmentation, the use of such polypeptides in screening assays, and kits comprising these reagents.</p> <b>(57) Abrégé</b> <p>La présente invention concerne des polypeptides humains isolés et purifiés à fonction kinase, les acides nucléiques codant ces polypeptides, des procédés de production de formes recombinées de ces polypeptides, des anticorps produits contre ces polypeptides, et des fragments de peptides obtenus à partir de ces polypeptides. Par ailleurs, cette invention concerne l'utilisation de ces polypeptides et des peptides fragmentés dans des réactions de phosphorylation et comme étalons de masse moléculaire, l'utilisation de ces polypeptides et des peptides fragmentés comme régulateurs de la fragmentation des peptides, l'utilisation de ces polypeptides dans des essais de dépistage, et enfin, des kits renfermant ces réactifs.</p>		

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 7 :</b> <b>C12N 15/55, 9/12, C07K 16/40, G01N 33/50</b>		<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/08180</b> <b>(43) International Publication Date:</b> 17 February 2000 (17.02.00)
<b>(21) International Application Number:</b> PCT/US99/17630 <b>(22) International Filing Date:</b> 3 August 1999 (03.08.99)  <b>(30) Priority Data:</b> 60/095,270 4 August 1998 (04.08.98) US 60/099,972 11 September 1998 (11.09.98) US  <b>(71) Applicant (for all designated States except US):</b> IMMUNEX CORPORATION [US/US]; 51 University Street, Seattle, WA 98101 (US).  <b>(72) Inventors; and</b> <b>(73) Inventors/Applicants (for US only):</b> VIRCA, G., Duke [US/US]; 16690 S.E. 50th Place, Bellevue, WA 98006 (US). BIRD, Timothy, A. [GB/US]; 10804 Olallie Drive N.E., Bainbridge Island, WA 98110 (US). ANDERSON, Dirk, M. [US/US]; 3616 N.W. 64th Street, Seattle, WA 98107 (US). MARKEN, John, S. [US/US]; 11804 9th Avenue S.W., Seattle, WA 98146 (US).  <b>(74) Agent:</b> JONES, Simone, L.; Immunex Corporation, Law Dept., 51 University Street, Seattle, WA 98101 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title:</b> HUMAN cDNAs ENCODING POLYPEPTIDES HAVING KINASE FUNCTIONS			
<b>(57) Abstract</b>  The invention is directed to purified and isolated human polypeptides having kinase function, the nucleic acids encoding such polypeptides, processes for production of recombinant forms of such polypeptides, antibodies generated against these polypeptides, fragmented peptides derived from these polypeptides, the use of such polypeptides and fragmented peptides in phosphorylation reactions and as molecular weight markers, the use of such polypeptides and fragmented peptides as controls for peptide fragmentation, the use of such polypeptides in screening assays, and kits comprising these reagents.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## D scription

5

10

15

20

25

30

35

40

45

50

55

## HUMAN cDNAs ENCODING POLYPEPTIDES HAVING KINASE FUNCTIONS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is related to the U.S. provisional application serial number 60/095,270, filed on August 4, 1998, and U.S. provisional application serial number 60/099,972, filed September 11, 1998, and bearing the same title and listing the same inventors.

FIELD OF THE INVENTION

The invention is directed to purified and isolated human polypeptides having kinase function, the nucleic acids encoding such polypeptides, processes for production of recombinant forms of such polypeptides, antibodies generated against these polypeptides, fragmented peptides derived from these polypeptides, the use of such polypeptides and fragmented peptides in phosphorylation reactions and as molecular weight markers, the use of such polypeptides and fragmented peptides as controls for peptide fragmentation, the use of such polypeptides in screening assays, and kits comprising these reagents.

BACKGROUND OF THE INVENTION

The eukaryotic protein kinases make up a large and rapidly expanding family of proteins related on the basis of homologous catalytic domains. Spurred by the development of gene cloning and sequencing methodologies, distinct protein kinase genes have been identified from a wide selection of invertebrates and lower eukaryotes, including *Drosophila*, *Caenorhabditis elegans*, *Aplysia*, *Hydra*, *Dictyostelium*, and budding (*Saccharomyces cerevisiae*) and fission (*Schizosaccharomyces pombe*) yeast. Homologous genes have also been identified in higher plants. Protein kinases, however, are not limited to the eukaryotes. Enzyme activities have been well documented in prokaryotes, but the prokaryotic protein kinase genes are not obviously homologous to those of the eukaryotes. Because protein kinases are useful biochemical reagents, there is a need in the art for the continued discovery of unique members of the protein kinase family.

In addition, the discovery and identification of proteins are at the forefront of modern molecular biology and biochemistry. The identification of the primary structure, or sequence, of a sample protein is the culmination of an arduous process of

5 experimentation. In order to identify an unknown sample protein, the investigator can  
rely upon comparison of the unknown sample protein to known peptides using a variety  
of techniques known to those skilled in the art. For instance, proteins are routinely  
10 analyzed using techniques such as electrophoresis, sedimentation, chromatography, and  
5 mass spectrometry.

Comparison of an unknown protein sample to polypeptides of known molecular  
weight allows a determination of the apparent molecular weight of the unknown protein  
15 sample (T.D. Brock and M.T. Madigan, *Biology of Microorganisms* 76-77 (Prentice Hall,  
6d ed. 1991)). Protein molecular weight standards are commercially available to assist in  
10 the estimation of molecular weights of unknown protein samples (New England Biolabs  
Inc. Catalog:130-131, 1995; J. L. Hartley, U.S. Patent No. 5,449,758). However, the  
20 molecular weight standards may not correspond closely enough in size to the unknown  
sample protein to allow an accurate estimation of apparent molecular weight.

The difficulty in estimation of molecular weight is compounded in the case of  
25 proteins that are subjected to fragmentation by chemical or enzymatic means (A.L.  
Lehninger, *Biochemistry* 106-108 (Worth Books, 2d ed. 1981)). Chemical fragmentation  
can be achieved by incubation of a protein with a chemical, such as cyanogen bromide,  
which leads to cleavage of the peptide bond on the carboxyl side of methionine residues  
30 (E. Gross, *Methods in Enz.* 11:238-255, 1967). Enzymatic fragmentation of a protein can  
20 be achieved by incubation of a protein with a protease that cleaves at multiple amino acid  
residues (D. W. Cleveland et al., *J. Biol. Chem.* 252:1102-1106, 1977). Enzymatic  
fragmentation of a protein can also be achieved by incubation of a protein with a protease,  
35 such as *Achromobacter* protease I (F. Sakiyama and A. Nakata, U.S. Patent No.  
5,248,599; T. Masaki et al., *Biochim. Biophys. Acta* 660:44-50, 1981; T. Masaki et al.,  
25 *Biochim. Biophys. Acta* 660:51-55, 1981), which leads to cleavage of the peptide bond on  
the carboxyl side of lysine residues. The molecular weights of the fragmented peptides  
40 can cover a large range of molecular weights and the peptides can be numerous.  
Variations in the degree of fragmentation can also be accomplished (D. W. Cleveland et  
al., *J. Biol. Chem.* 252:1102-1106, 1977).

45 30 The unique nature of the composition of a protein with regard to its specific amino  
acid constituents results in a unique positioning of cleavage sites within the protein.  
Specific fragmentation of a protein by chemical or enzymatic cleavage results in a unique  
"peptide fingerprint" (D. W. Cleveland et al., *J. Biol. Chem.* 252:1102-1106, 1977; M.  
50 Brown et al., *J. Gen. Virol.* 50:309-316, 1980). Consequently, cleavage at specific sites

5 results in reproducible fragmentation of a given protein into peptides of precise molecular weights. Furthermore, these peptides possess unique charge characteristics that determine the isoelectric pH of the peptide. These unique characteristics can be exploited using a variety of electrophoretic and other techniques (T.D. Brock and M.T. Madigan, *Biology of Microorganisms* 76-77 (Prentice Hall, 6d ed. 1991)).

10 When a peptide fingerprint of an unknown protein is obtained, this can be compared to a database of known proteins to assist in the identification of the unknown protein (W.J. Henzel et al., *Proc. Natl. Acad. Sci. USA* 90:5011-5015, 1993; B. Thiede et al., *Electrophoresis* 1996, 17:588-599, 1996). A variety of computer software programs are accessible via the Internet to the skilled artisan for the facilitation of such comparisons, such as MultiIdent (Internet site: [www.expasy.ch/sprot/multiident.html](http://www.expasy.ch/sprot/multiident.html)), PeptideSearch (Internet site: [www.mann.embl-heidelberg.de...deSearch/FR\\_PeptideSearchForm.html](http://www.mann.embl-heidelberg.de...deSearch/FR_PeptideSearchForm.html)), and ProFound (Internet site: [www.chait-sgi.rockefeller.edu/cgi-bin/prot-id-frag.html](http://www.chait-sgi.rockefeller.edu/cgi-bin/prot-id-frag.html)). These programs allow the user to specify the cleavage agent and the molecular weights of the fragmented peptides within a designated tolerance. The programs compare these molecular weights to protein databases to assist in the elucidation of the identity of the sample protein. Accurate information concerning the number of fragmented peptides and the precise molecular weight of those peptides is required for accurate identification. Therefore, increasing the accuracy in the determination of the number of fragmented peptides and the precise molecular weight of those peptides should result in enhanced success in the identification of unknown proteins.

35 Fragmentation of proteins is further employed for the production of fragments for amino acid composition analysis and protein sequencing (P. Matsudaira, *J. Biol. Chem.* 262:10035-10038, 1987; C. Eckerskorn et al., *Electrophoresis* 1988, 9:830-838, 1988), particularly the production of fragments from proteins with a "blocked" N-terminus. In addition, fragmentation of proteins can be used in the preparation of peptides for mass spectrometry (W.J. Henzel et al., *Proc. Natl. Acad. Sci. USA* 90:5011-5015, 1993; B. Thiede et al., *Electrophoresis* 1996, 17:588-599, 1996), for immunization, for affinity selection (R. A. Brown, U.S. Patent No. 5,151,412), for determination of modification sites (e.g. phosphorylation), for generation of active biological compounds (T.D. Brock and M.T. Madigan, *Biology of Microorganisms* 300-301 (Prentice Hall, 6d ed. 1991)), and for differentiation of homologous proteins (M. Brown et al., *J. Gen. Virol.* 50:309-316, 1980).

In view of the continuing interest in protein research and the elucidation of protein structure and properties, there exists a need in the art for polypeptides having kinase function or suitable for use in peptide fragmentation studies and in molecular weight measurements.

#### **SUMMARY OF THE INVENTION**

The invention aids in fulfilling these needs in the art. The invention encompasses an isolated human nucleic acid molecule comprising the DNA sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 13, or 15 and an isolated human nucleic acid molecule encoding the amino acid sequence of SEQ ID NO:7, 8, 9, 10, 11, 12, 14, or 16. The invention also encompasses nucleic acid molecules complementary to these sequences. As such, the invention includes double-stranded nucleic acid molecules comprising the DNA sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 13, or 15 and isolated nucleic acid molecules encoding the amino acid sequence of SEQ ID NO:7, 8, 9, 10, 11, 12, 14 or 16. Both single-stranded and double-stranded RNA and DNA nucleic acid molecules are encompassed by the invention. These molecules can be used to detect both single-stranded and double-stranded RNA and DNA variants encompassed by the invention. A double-stranded DNA probe allows the detection of nucleic acid molecules equivalent to either strand of the nucleic acid molecule. Isolated nucleic acid molecules that hybridize to a denatured, double-stranded DNA comprising the DNA sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 13, or 15, or an isolated nucleic acid molecule encoding the amino acid sequence of SEQ ID NO:7, 8, 9, 10, 11, 12, 14, or 16 are within the invention. A preferred set of hybridization conditions are those of moderate stringency: in 50% formamide and 6XSSC, at 42°C with washing conditions of 60°C, 0.5XSSC, 0.1% SDS.

The invention further encompasses isolated nucleic acid molecules derived by *in vitro* mutagenesis from SEQ ID NO:1, 2, 3, 4, 5, 6, 13, or 15. *In vitro* mutagenesis would include numerous techniques known in the art including, but not limited to, site-directed mutagenesis, random mutagenesis, and *in vitro* nucleic acid synthesis. The invention also encompasses isolated nucleic acid molecules degenerate from SEQ ID NO:1, 2, 3, 4, 5, or 6 (and the resulting amino acid sequence) as a result of the genetic code, isolated nucleic acid molecules that are allelic variants of human DNA of the invention, or a species homolog of DNA of the invention. The invention also encompasses recombinant vectors that direct the expression of these nucleic acid molecules and host cells transformed or transfected with these vectors. In addition, the invention encompasses methods of using



the nucleic acid noted above in assays to identify chromosomes, map human genes, and study tumors.

The invention also encompasses isolated polypeptides encoded by these nucleic acid molecules, including isolated polypeptides having a molecular weights as determined by SDS-PAGE, isolated polypeptides in non-glycosylated form, and fragments thereof. The invention further includes synthetic polypeptides encoded by these nucleic acid molecules. Peptides and fragments of these polypeptides, however derived, are also part of the invention and may be produced by any standard means, from chemical, enzymatic, recombinant, or synthetic methods. Isolated polyclonal or monoclonal antibodies that bind to these polypeptides are encompassed by the invention. The invention further encompasses methods for the production of polypeptides having kinase functions including culturing a host cell under conditions promoting expression and recovering the polypeptide from the culture medium. Especially, the expression of polypeptides having kinase functions in bacteria, yeast, plant, insect, and animal cells is encompassed by the invention.

In general, the polypeptides of the invention having kinase function can be used to phosphorylate target proteins and to radiolabeled target proteins with  $^{32}\text{P}$ . In addition, the polypeptides of the invention having kinase function can be used to identify proteins having a phosphate activity.

In addition, assays utilizing polypeptides having kinase functions to screen for potential inhibitors of activity associated with polypeptide counter-structure molecules, and methods of using polypeptides having kinase functions as therapeutic agents for the treatment of diseases mediated by polypeptide counter-structure molecules are encompassed by the invention. Methods of using polypeptides having kinase functions in the design of inhibitors thereof are also an aspect of the invention. The invention further encompasses use of polypeptides of the invention to screen for agonists and antagonists.

The invention further encompasses the fragmented peptides produced from polypeptides of the invention by chemical or enzymatic treatment. In addition, the polypeptides of the invention and fragmented peptides thereof, wherein at least one of the sites necessary for fragmentation by chemical or enzymatic means has been mutated, are an aspect of the invention.

The invention further includes a method for using these polypeptides and fragmented peptides thereof as molecular weight markers that allow the estimation of the molecular weight of a protein or a fragmented protein sample. The invention also

5 encompasses a method for the visualization of the molecular weight markers of the invention thereof using electrophoresis. The invention further encompasses methods for using the polypeptides of the invention and fragmented peptides thereof as markers, which  
10 aid in the determination of the isoelectric point of a sample protein. The invention also encompasses methods for using polypeptides of the invention and fragmented peptides thereof as controls for establishing the extent of fragmentation of a protein sample.

15 Further encompassed by this invention are kits to aid the determination of molecular weights of a sample protein utilizing polypeptide molecular weight markers of the invention, fragmented peptides thereof, and forms of these polypeptide molecular  
20 weight markers, wherein at least one of the sites necessary for fragmentation by chemical or enzymatic means has been mutated.

### DETAILED DESCRIPTION OF THE INVENTION

25 The protein kinases are a large family of enzymes, many of which mediate the response of eukaryotic cells to external stimuli. In recent years, members of the protein kinase family have been discovered at an accelerated pace. The surge in the number of known protein kinases has been largely due to the advent of gene cloning and sequencing  
30 techniques. Amino acid sequences deduced from nucleotide sequences are considered to represent protein kinases if they include certain key residues that are highly conserved in the protein kinase catalytic domain. A cDNA encoding a human polypeptide has been isolated and is set forth in SEQ ID NO:1, 2, 3, 4, 5, 6, 13, or 15. This discovery of the  
35 cDNA encoding human polypeptides having kinase functions enables construction of expression vectors comprising nucleic acid sequences encoding polypeptides having kinase functions; host cells transfected or transformed with the expression vectors;  
25 biologically active human polypeptides having kinase functions, and molecular weight markers as isolated and purified proteins; and antibodies immunoreactive with  
40 polypeptides of the invention.

More particularly, the invention relates to certain nucleotide sequences. A  
45 "nucleotide sequence" refers to a polynucleotide molecule in the form of a separate fragment or as a component of a larger nucleic acid construct, that has been derived from DNA or RNA isolated at least once in substantially pure form (i.e., free of contaminating  
50 endogenous materials) and in a quantity or concentration enabling identification, manipulation, and recovery of its component nucleotide sequences by standard biochemical methods (such as those outlined in Sambrook et al., *Molecular Cloning: A*

Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989)). Such sequences are preferably provided and/or constructed in the form of an open reading frame uninterrupted by internal non-translated sequences, or introns, that are typically present in eukaryotic genes. Sequences of non-translated DNA can be present 5' or 3' from an open reading frame, where the same do not interfere with manipulation or expression of the coding region.

Particularly preferred nucleotide sequences of the invention include the following:

**NAME: HH0900-BF04 DNA**

**Nucleotide sequence:**

GTACGCCATGAAGGTGCTGCGCAAGGCGGCGTGGTGACGCGCCAAAGA  
CGCAAGAGCACACGCGCACCAGCGCTCGGTGCTGGAGCTGGTGCGCCAG  
GCGCCCTTCCTGGTCACGCTGCACTACGCTTCCAGACGGATGCCAAGCT  
GCACCTCATCCTGGACTATGTGAGCGGCGGG (SEQ ID NO:1);

**NAME: HH2040-BF04 DNA**

**Nucleotide sequence:**

CCCGAGAGGTGCCACATCAGACCGCTCCGACTTCGTGCGGGACTCGGCG  
GCCAGCCACCAGGCGGAGCCCGAGGCGTACGAGCGGCGCGTGTGCTTCCT  
GCTTCTGCAACTCTGCAACGGGCTGGAGCACCTGAAGGAGCACGGGATCA  
TCCACCGGACCTGTGCCTGGAGAACCTGTGCTGGTGCACTGCACCCTC  
CAGGCCGGCCCCGGGCCCCGCC (SEQ ID NO:2);

**Name: JJ503-KS DNA**

**nucleotide sequence**

CGGGCAGGGCTGGAGCTGGGCTGGGATCCCGAGCTCGGCAGCAGCGCAGCGGGCCGGCCACCTGCTGGTG  
CCTGGAGGCTCTGAGCCCCGGCGGCGCCCGGGCCACGCGGAACGACGGGGCGAGATGCGAGCCACCCCTCT  
GGCTGCTCCTGCGGGTTCCCTGTCCAGGAAGAAGCGGTTGGAGTTGGATGACAACTTAGATACCGAGCGTCC  
CGTCCAGAAACGAGCTCGAAGTGGGCCCCAGCCAGACTGCCCCCTGCCCTGTTGCCCCCTGAGCCACCTAC  
TGCTCCAGATCGTGCAACTGCTGTGGCCACTGCCTCCCGTCTTGGGCCCTATGTCTCCTGGAGCCCGAGGA  
GGGCGGGCGGGCTTACCAGGCCCTGCCTGCCCCACAGGCACTGAGTATACCTGCAAGGTGTACCCCGTCCA  
GGAAGCCCTGGCCGTGCTGGAGCCCTACGCGCGGCTGCCCCCGCACAAGCATGTGGCTCGGGCCACTGAGGT  
CCTGGCTGGTACCCAGCTCCTCTACGCCTTTTCACTCGGACCCATGGGGACATGCACAGCCTGGTGCGAAG  
CCGCCACCGTATCCCTGAGCCTGAGGCTGCCGTGCTCTTCCGCCAGATGGCCACCGCCCTGGCGCACTGTCA  
CCAGCACGGTCTGGTCTGCGTGATCTCAAGCTGTGTCGCTTTGTCTTCGCTGACCGTGAGAGGAAGAAGCT  
GGTGCTGGAGAACCCTGGAGGACTCCTGCGTGCTGACTGGGCCAGATGATTCCCTGTGGGACAAGCACGCGTG  
CCCAGCCTACGTGGGACCTGAGATACTCAGCTCACGGGCCTCATACTCGGGCAAGGCAGCCGATGTCTGGAG  
CCTGGGCGTGGCGCTCTTACCATGCTGGCCGGCCACTACCCCTTCCAGGACTCGGAGCCTGTCTGCTCTT  
CGGCAAGATCCGCCCGGGGCGCTACGCCTTGCTGCGAGGCTCTCGGCCCTGCCCCGCTGTCTGGTTCGCTG  
CCTCCTTCGTCGGGAGCCAGCTGAACGGCTCACAGCCACAGGCATCCTCTGCACCCTGGCTGCGACAGGA  
CCCGA (SEQ ID NO:3);

Name: QQ1249-BF04 DNA

Nucleotide sequence:

CAGCGAGAAGCCGACATGCATCGCCTCTTCAATCACCCCAACATCCTTCG  
5 CCTCGTGGCTTACTGTCTGAGGGAACGGGGTGCTAAGCATGAGGCCTGGC  
TGCTGCTACCATTCCTTCAAGAGAGGTACGCTGTGGAATGAGATAGAAAGG  
CTGAAGGACAAAGGCAACTTCCTGACCGAGGATCAAATCCTTTGGCTGCT  
GCTGGGGATCTGCAGAGGCCTTGAGGCCATTTCATGCCAAGGGTTATGCCT  
10 ACAGAGACTTGAAGCCCAACAATATATTGCTTGGAGATGAGGGGCAGCCA  
GTTTAAATGGACTTGGGTTCATGAATCAAGCATGCATCCATGTGGAGGG  
15 CTCCCGCCAGGCTCTGACCTGCAGGACTGGGCAGCCC (SEQ ID NO: 4);

Name: QQ3351-BF04 DNA

Nucleotide sequence:

15 ATGCTAACTAGTTTAAACAGATCTTGAACGAGACGACCTGCTGTGGAAGAGCGAGCTTTTGGAACTGTGC  
ACGGGACAGATTGGACGCACACCCCTCGGGAGGCGCGAAGGCATGGAAAATTTGAAGCATATTATCACCTT  
GGCCAGGTCATCCACAAACGGTGTGAAGAGATGAAATACTGCAAGAAACAGTGCCGGCGCTGGGCCACCGC  
GCTCCTCGGCCTGATCAAGCCTCTGGAGATGCTCCAGGACCAAGGAAAGAGAGCGTGCCTCTGAGAAGTTA  
20 ACCACAGCCATGAACCCTTCAAGGCTGCCCTGGAGGAGGCTAATGGGGAGATAGAAAAGTTCAGCAATAGA  
TCCAATATCTGCAGGTTTCTAACAGCAAGCCAGGACAAAATACTCTTCAAGGACGTGAACAGGAAGCTGAGT  
25 GATGTCTGGAAGGAGCTCTCGCTGTTACTTCAGGTTGAGCAACGCATGCCTGTTTCACCCATAAGCCAAGGA  
GCGTCTTGGGCACAGGAAGATCAGCAGGATGCAGACGAAGACAGGCGAGCTTTCCAGATGCTAAGAAGAGAT  
AATGAAAAAATAGAAGCTTCACTGAGACGATTAGAAAATCAACATGAAAGAAATCAAGGAAACTTTGAGGCAG  
30 TATTTACCACAAAATGCATGCAGGAGATCCCGCAAGAGCAAAATCAAGGAGATCAAGAAGGAGCAGCTTTCA  
GGATCCCCGTGGATTCTGCTAAGGGAAAATGAAGTCAGCACACTTTATAAAGGAGAAATACCACAGAGCTCCA  
25 GTGGCCATAAAAAGTATTCAAAAAACTCCAGGCTGGCAGCATTGCAATAGTGAAGCAGACTTTCAATAAGGAG  
ATCAAAACCATGAAGAAATTCGAATCTCCCAACATCCTGCGTATATTTGGGATTGCAATGATGAACAGTG  
ACTCCGCCCTCAATTCTCCATTGTCTGAGTACTGTGAACTCGGGACCTTGAGGGAGCTGTTGGATAGGGAA  
30 AAAGACCTCACACTTGGCAAGCGCATGGTCTAGTCTTGGGGCAGCCCGAGGCTTATACCGCTACACCAT  
TCAGAAGCACCTGAACCTCCACGGAAAAATCAGAAGCTCAAACCTTCTGGTAACCAAGGCTACCAAGTGAAG  
CTTGAGGATTTGAGTTGAGGAAAAACACAGACTTCCATGAGTTTGGGAACTACGAGAGAAAAGACAGACAGA  
GTCAAATCTACAGCATATCTCTACCTCAGGAACTGGAAGATGTATTTTATCAATATGATGTAAAGTCTGAA  
40 ATATACAGCTTTGGAATCGTCTCTGGGAAATCGCCACTGGAGATATCCCGTTTCAAGGCTGTAATTCTGAG  
AAGATCCGCAAGCTGGTGGCTGTGAAGCGGCAGCAGGAGCCACTGGGTGAAGACTGCCCTTCAGAGCTGCGG  
35 GAGATCATTGATGAGTGCCGGGCAGCAGGTCGCTCTGTTCCAAGATCTGTAGCGGCCGCCCGGCCGCTCGAC  
GTTTAAACGCGTGGCCCTCGAGAGGTTTTCGATCCGTCGAT (SEQ ID NO:5), and

Name: SS1771

Nucleotide sequence:

CTTCCCCTG GACGTGGAGT ACGGAGGCCC AGACCGGAGG TGCCCCCTC  
CGCCCTACCC GAAGCACCTG CTGCTGCGCA GCAAGTCGGA GCAGTACGAC  
45 CTGGACAGCC TGTGCGCAGG CATGGAGCAG AGCCTCCGTG CGGGCCCCAA  
CGAGCCCGAG GCGGCGGACA AGAGCCGCAA AAGCGCCAAG GGGGACAAAG  
GCGGAAAGGA TAAAAAGCAG ATTCAAGACT CTCCCGTTCC CGTCCGCAAA  
AACAGCAGAG ACGAAGAGAA GAGAGAGTCA CGCATCAAGA GCTACTCGCC  
ATACGCCTTT AAGTTCTTCA TGGAGCAGCA CGTGGAGAAT GTCATCAAAA  
50 CCTACCAGCA GAAGGTTAAC CGGAGGCTGC AGCTGGAGCA AGAAATGGCC  
AAAGCTGGAC TCTGTGAAGC TGAGCAGGAG CAGATGCGGA AGATCCTCTA  
CCAGAAAGAG TCTAATTACA ACAGGTTAAA GAGGGCCAAG ATGGACAAGT  
55 CTATGTTTGT CAAGATCAAA ACCCTGGGGA TCGGTGCCTT TGGAGAAGTG

5  
10  
15  
20  
20

TGCCCTTGCTT GTAAGGTGGA CACTCACGCC CTGTACGCCA TGAAGACCCT  
AAGGAAAAAG GATGTCTCTGA ACCGGAATCA GGTGGCCAC GTCAAGGCCG  
AGAGGGACAT CCTGGCCGAG GCAGACAATG AGTGGGTGGT CAAACTCTAC  
TACTCCTTCC AAGACAAAGA CAGCCTGTAC TTTGTGATGG ACTACATCCC  
5 TGGTGGGGAC ATGATGAGCC TGCTGATCCG GATGGAGGTC TTCCCTGAGC  
ACCTGGCCCG GTTCTACATC GCAGAGCTGA CTTTGGCCAT TGAGAGTGTC  
10 CACAAGATGG GCTTCATCCA CCGAGACATC AAGCCTGATA ACATTTTGAT  
AGATCTGGAT GGTACATTA AACTCACAGA TTTCCGGCTC TGCCTGGGT  
TCAGGTGGAC TCACAATTCC AAATATTACC AGAAAGGGAG CCATGTCAGA  
10 CAGGACAGCA TGGAGCCAG CGACCTCTGG GATGATGTGT CTAAGTGTG  
GTGTGGGGAC AGGCTGAAGA CCCTAGAGCA GAGGGCCGG AAGCAGCACC  
AGAGGTGCC TGGCACATTCA CTGGTGGGA CTCCAACTA CATCGCACC  
GAGGTGCTCC TCCGCAAGG GTACACTCAA CTCTGTGACT GGTGGAGTGT  
15 TGGAGTGATT CTCTTCGAGA TGCTGGTGGG GCAGCCGCC TTTTGGCAC  
15 CTACTCCAC AGAAACCCAG CTGAAGGTGA TCAACTGGGA GAACACGCTC  
CACATTCCAG CCCAGGTGAA GCTGAGCCCT GAGGCCAGG ACCTCATCAC  
CAAGCTGTG TGCTCCGAG ACCACGCCCT GGGGCGGAAT GGGGCCGATG  
ACCTGAAGC CCACCCCTC TTCAGCGCCA TTGACTTCTC CAGTGACATC  
CGGAAGCATC CAGCCCCCTA CGTCCCACC ATCAGCCACC CCATGGAG  
20 (SEQ ID NO:6).

Name: SS1771A

25  
30  
35  
40  
45  
50  
55

**Nucleotide sequence:**  
TCCCCTGGACGTGGAGTACGGAGGCCAGACCGGAGGTGCCCGCCTCCGCCCTACCCGAAGCACCTGCTGC  
TGCCGAGCAAGTCGGAGCAGTACGACCTGGACAGCCTGTGCGCAGGCATGGAGCAGAGCCTCCGTGCGGGCC  
CCACAGAGCCCGAGGGCGGCACAGAGCCGCAAAAGCGCCAAAGGGGACAAAGCCGAAAGGATAAAAAGC  
AGATTACAGACCTCTCCCGTTCCCGTCCCGCAAAACAGCAGAGACGAAGAGAGAGAGTACGCATCAAGA  
30 GCTACTCGCCATACGCCCTTTAAGTTCTTCATGGAGCAGCACGTGGAGAATGTCATCAAAACCTACCAGCAGA  
AGGTTAAACCGGAGGCTGCAGCTGGAGCAAGAAATGGCCAAAGCTGGACTCTGTGAAGCTGAGCAGGAGCAGA  
TGCGGAAGATCCTCTACCAGAAAGAGTCTAATTACAACAGGTTAAAGAGGGCCAAAGATGGACAAGTCTATGT  
35 TTGTCAAGATCAAAACCTGGGGATCGGTGCCCTTTGGAGAAGTGTGCCTTGCTTGTAAGGTGGACACTCACG  
CCCTGTACGCCATGAAGACCCTAAGGAAAAAGGATGTCTGAACCGGAATCAGGTGGCCACGTCAAGGCCG  
AGAGGGACATCCTGGCCGAGGCAGACAATGAGTGGTGGTCAAACCTACTACTCCTTCCAAGACAAGACA  
40 TCTGCTACTGGGTTTCAGGTGGACTCACAAATTCCAAATATTACCAGAAAGGGAGCCATGTCAGACAGGACAGCA  
TGGAGCCCGACGACCTCTGGGATGATGTGTCTAACTGTCCGGTGTGGGACAGGCTGAAGACCCCTAGAGCAGA  
GGGCGCGGAAGCAGCACCAGAGGTGCCCTGGCACATTCACCTGGTGGGACTCCAACTACATCGCACCAGG  
TGCTCCTCCGCAAGGGTACACTCAACTCTGTGACTGGTGGAGTGTGGAGTGATCTCTTCGAGATGCTGG  
45 TGGGCGAGCCGCCCTTTTGGCACTACTCCACAGAAACCCAGCTGAAGGTGATCAACTGGGAGAACACGC  
TCCACATTCAGCCAGGTGAAGCTGAGCCCTGAGGCCAGGGACCTCATCACCAGCTGTGCTGCTCCGCAG  
ACCACCGCTGGGGCGGAATGGGGCCGATGACCTGAAGGCCACCCCTTCTTCAGCGCCATTGACTTCTCCA  
50 GTGACATCCGGAAGCATCCAGCCCTACGTTCCCAACATCAGCCACCCCATGGACACCTCGAATTTCGACC  
CCGTAGATGAAGAAAGCCCTTGGAAAGATGCCAGCGAAGGTAGCACCAGGCTGGGACACACTACCTCGC  
CCAATAACAAGCATCCTGAGCACGCATTTTACGAATTCACCTTCCGAAGGTCTTTGATGACAATGGCTACC  
CCTTTCGATGCCAAAGCCTTCAGGAGCAGAAGCTTCACAGGCTGAGAGCTCAGATTTAGAAAGCTCTGATC  
55 TGGTGGATCAGACTGAAGGCTGCCAGCCTGTGTACGTGTAGATGGGGCCAGGCACCCCACTCGCTGC  
CTCCAGGTCAGGGTCCCGAGCCGGTCCCTCACAGGCCAATAGGGAAGCCGAGGGCTGTTTTGTTTTAAA  
TTAGTCCGTCGATTACTTCACTTGAATCTTGCTCTTACCAGAAACCCAAACAGGACACTTTTGAAAAAC  
AGCGGTGCCCGGAATTC (SEQ ID NO:15).

With the continued increase in the number of known eukaryotic protein kinases, a  
suitable classification scheme is one based on comparing catalytic-domain sequences. It  
follows that protein kinases with similar catalytic domains will tend also to have similar

enzymatic and regulatory properties. The nucleotide sequences of the invention encode, respectively, the following polypeptides having kinase function:

**NAME: HH0900-BF04 Polypeptide**

**Translation in relevant reading frame (3'-5' Frame 3):**

YAMKVLKKAALVQRAKTQEHTRTERSVLELVRQAPFLVTLHYAFQTDAKLHLILDY  
VSGG (SEQ ID NO:7);

The above sequence is consistent with the consensus sequence of subdomains II through V of the eukaryotic protein kinase superfamily. This sequence is the sequence of a polypeptide having kinase activity encoded by the nucleotide sequence of SEQ ID NO:1.

**NAME: HH2040-BF04 Polypeptide**

**Translation in relevant reading frame (5'-3' Frame 2):**

REVPHQTASDFVRDSAASHQAEPEAYERRVCFLLLQLCNGLE  
HLKEHGI IHRDLCLENLLLVHCTLQAGPGPA (SEQ ID NO:8);

The above sequence is consistent with the consensus sequence of subdomains V, VIA and VIB of the eukaryotic protein kinase superfamily. This sequence is the sequence of a polypeptide having kinase activity encoded by the nucleotide sequence of SEQ ID NO:2.

**Name: JJ503-KS Polypeptide**

**Translation in relevant reading frame (5'-3' frame 1):**

GQGWSWAGIPSSAAQ RAGPPAGALEALSPGGARAHAEERRGEMRATPLAA  
PAGSLSRKKRLELDDNLDTERPVQKRARSGPQPRLPCLLPLSPPTAPDR  
ATAVATASRLGPYVLEPEEGGRAYQALHCPTGTETCKVYPVQEALAVL  
EPYARLPPHKHVARPTEVLAGTQLLYAFFTRTHGDMHSLVRSRHRIPEPE  
AAVLFRQMATALAHCHQHGLVLRDLKLCRFVFADRERKKLVLENLEDCV  
LTGPDDSLWDKHACPAYVGPEILSSRASYSKGAADVWSLGVALFTMLAGH  
YPFQDSEPVLLFGKIRRGAYALPAGLSAPARCLVRCLLRREPAERLTATG  
ILLHPWLRQD (SEQ ID NO:9);

The sequence is consistent with kinase domains VIA through XI, though the homology is imperfect. This sequence is the sequence of a polypeptide having kinase activity encoded by the nucleotide sequence of SEQ ID NO:3.

**Name:** QQ1249-BF04 Polypeptide

**Translation in relevant reading frame (5'-3' frame 3)**

QREADMHRLFNHPNILLRLVAYCLRERGAKEAWLLPFFKRGTLWNEIER  
LKDKGNFLTQILWLLLGICRGLEATHAKGYAYRDLKPTNILLGDEGQP  
VLMDLGS MNQACIHVEGSRQALTLQDWAAQRCTISYRAPXLFVSQS  
(SEQ ID NO:10);

The above sequence is consistent with the consensus sequence of subdomains III - VIII of the eukaryotic protein kinase superfamily. This sequence is the sequence of a polypeptide having kinase activity encoded by the nucleotide sequence of SEQ ID NO:4.

**Name:** QQ3351-BF04 Polypeptide

**Translation in relevant frame (5'-3' frame 1)**

MLTSLNRSWNETTCCGRASFLELCTGGIGRTPLGRREGMENLKHIIITLGQVIHKRCEEMKYCKKQCRRRLGHR  
VLGLIKPLEMLQDQGRSVPSEKLTAMNRFKAAL EEANGEIEKFSNRSNICRFLTASQDKILFKDVNRKLS  
DVWKELSLLLQVEQRMVSPISQASWAQEDQDADDDRRAFQMLRRDNEKIEASLRRLINMKEIKETLRQ  
YLPKCMQEI PQEQIKEIKKEQLSGSPWILLRENEVSTLYKGEYHRAPVAIKVFKKLQAGSIATVRQTFNKE  
IKTMKKFESPNILRIFGICIDETVTPPQFSIVMEYCELGTLRELLDREKDLTLGKRMVLVLGAARGLYRLHH  
SEAPELHGKIRSSNFLTQGYQVLAGFELRKTQTSMSLGTTREKTRVKSTAYLSPQELEDDVFYQYDVKSE  
IYSGIVLWEIATGDI PFQGCNSEKIRKLVAVKRQEQEPLGEDCPSSELREI IDECRAAGRLVPRSVAAARA V  
(SEQ ID NO:11)

The above sequence is believed to be full-length. However, the initial methionine was not present in the clone sequenced but subsequently was added by PCR. Therefore, the natural sequence may comprise Leu-2 through the Val-505. This sequence is the sequence of a polypeptide having kinase activity encoded by the nucleotide sequence of SEQ ID NO:5.

**Name:** SS1771

**Translation in relevant frame (3'-5' frame 3)**

FPLDVEYGGPDRRCPPFPYPKHLLLRSKSEQYDLDLSCAGMEQSLRAGPN  
EPEGGDKSRKSAKGDKGKDKKQIQTSVPVVRKNSRDEEKRESRIKSYSP  
YAFKFFMEQHVENVIKTYQQKVNRRQLLEQEMAKAGLCEAEQEQMRKILY  
QKESNYNRLKRAKMDKSMFVKIKTLGIGAFGEVCLACKVDTHALYAMKTL  
RKDKVLNRRNQVAHVKAERDILAEADNEWVVKLYYSFQDKDSL YFVMDYIP

GGDMSLLIRMEVFPPEHLARFYIAELTLAIESVHKMGFIHRDIKPDNILI  
 DLDGHIKLTDFGLCTGFRWTHNSKYYQKGSVHRQDSMEPSDLWDDVSNCR  
 CGDRLKLTLEQRARKQHQRCLAHSLVGTPTYIAPEVLLRKGYTQLCDWWSV  
 GVILFEMLVGQPPFLAPTPTETQLKVINWENTLHI PAQVKLSPEARDLIT  
 KLCCSADHRLGRNGADDLKAHPFFSAIDFSSDIRKHPAPYVPTISHPME  
 (SEQ ID NO:12).

This sequence is the sequence of a polypeptide having kinase activity encoded by the  
 nucleotide sequence of SEQ ID NO:6. The sequence may also comprise Pro-2 through  
 Glu-499.

Name: SS1771A

#### Translation in relevant frame (5'-3' frame 3)

PLDVEYGGPDRRCPPPPYPKHLRLRSKSEQYDLDLSLCAGMEQSLRAGPNEPEGGDKSRKSAKGDKGKDKKQ  
 IQTSPVVPVRKNSRDEEKRESRIKSYSPYAFKFFMEQHVENVIKTYQQKVNRLQLEQEMAKAGLCEAEQEQM  
 RKILYQKESNYNRLKRAKMDKSMFVKIKTLGIGAFGEVCLACKVDTHALYAMKTLRKKDVLNRNQVAHVKA  
 RDILAEADNEWVVKLYYSFQDKDSLYFVMDYIPGGDMSLLIRMEVFPPEHLARFYIAELTLAIESVHKMGFI  
 HRDIKPDNILIDL DGHKLTDFGLCTGFRWTHNSKYYQKGSVHRQDSMEPSDLWDDVSNCRCGDRLKLTLEQR  
 ARKQHQRCLAHSLVGTPTYIAPEVLLRKGYTQLCDWWSVGVILFEMLVGQPPFLAPTPTETQLKVINWENTL  
 HI PAQVKLSPEARDLITKLCCSADHRLGRNGADDLKAHPFFSAIDFSSDIRKHPAPYVPTISHPMDTSNFD  
 P VDEESPNWDASEGSTKAWDTLTSNNKHPEHAFYEFTFRFFDDNGYPFRCPKPSGAEASQAESSDLESSDL  
 VDQTEGCQPVYV (SEQ ID NO:16)

This sequence is the sequence of a polypeptide having kinase activity encoded by the  
 nucleotide sequence of SEQ ID NO:1.

The invention also includes truncated forms of the nucleic acids and polypeptides  
 of the invention. In a preferred embodiment, the invention includes a truncated form of  
 QQ3351-BF04, as follows:

#### Nucleotide sequence:

CTTGCAGGATTGAGTTGAGGAAAAACAGACTTCCATGAGTTTGGGAACACGAGAGAAAAGACAGACAGA  
 GTCAAATCTACAGCATATCTCTCACCTCAGGAACAGGAAGATGTATTTATCAATATGATGTAAAGCTGAA  
 ATATACAGCTTTGGAATCGTCTCTGGGAAATCGCCACTGGAGATATCCCGTTTCAAGGCTGTAATTCTGAG  
 AAGATCCGCAAGCTGGTGGCTGTGAAGCGGCAGCAGGAGCCACTGGGTGAAGACTGCCCTTCAGAGCTGCGG  
 GAGATCATTTGATGAGTGCCGGGCCATGATCCCTCTGTGCGGCCCTCTGTGGATGAAATCTTAAAGAAACTC  
 TCCACCTTTTCTAAG (SEQ ID NO:13)

#### Translation in relevant frame (5'3' frame 1):

LAGFELRKTQTSMSLGTREKTRVKSTAYLSPQLEDFYQYDVKSEIYS  
 FGIVLWEIATGDI PFQGCNSEKIRKLVAVKRQOEPLGEDCPSELREIIEC  
 RAHDPSVRPSVDEILKKLSTFSK (SEQ ID NO:14)

This sequence is the sequence of a polypeptide having kinase activity encoded by the  
 nucleotide sequence of SEQ ID NO:13.



5 The polypeptides of the invention are useful for characterizing cell and tissue expression, understanding their roles in development or hormonal response, and identifying regulatory molecules and physiologically relevant protein substrates.

10 As used herein, the term "polypeptides of the invention" refers to a genus of polypeptides that further encompasses proteins having the amino acid sequence of SEQ ID NO: 7, 8, 9, 10, 11, 12, 14, or 16 as well as those proteins having a high degree of similarity (at least 90% homology) with such amino acid sequences and which proteins are biologically active. In addition, polypeptides of the invention refers to the gene products of the nucleotides of SEQ ID NO:1, 2, 3, 4, 5, 6, 13 or 15.

15 The isolated and purified polypeptides of the invention have molecular weights of approximately 6883 (HH0900-BF04); 8168 (HH2040-BF04); 39,284 (JJ503-KS); 16,718 (QQ1249-BF04); 58,001 (QQ3351-BF04); 57,381 (SS1771), and 67,331 (SS1771A) Daltons in the absence of glycosylation. It is understood that the molecular weight of these polypeptides can be varied by fusing additional peptide sequences to both the amino and carboxyl terminal ends of polypeptides of the invention. Fusions of additional peptide sequences at the amino and carboxyl terminal ends of polypeptides of the invention can be used to enhance expression of these polypeptides or aid in the purification of the protein.

20 It is understood that fusions of additional peptide sequences at the amino and carboxyl terminal ends of polypeptides of the invention will alter some, but usually not all, of the fragmented peptides of the polypeptides generated by enzymatic or chemical treatment.

25 It is understood that mutations can be introduced into polypeptides of the invention using routine and known techniques of molecular biology. It is further understood that a mutation can be designed so as to eliminate a site of proteolytic cleavage by a specific enzyme or a site of cleavage by a specific chemically induced fragmentation procedure. It is also understood that the elimination of the site will alter the peptide fingerprint of polypeptides of the invention upon fragmentation with the specific enzyme or chemical procedure.

30 The term "isolated and purified" as used herein, means that the polypeptides or fragments of the invention are essentially free of association with other proteins or polypeptides, for example, as a purification product of recombinant host cell culture or as a purified product from a non-recombinant source. The term "substantially purified" as used herein, refers to a mixture that contains polypeptides or fragments of the invention

5 and is essentially free of association with other proteins or polypeptides, but for the presence of known proteins that can be removed using a specific antibody, and which substantially purified polypeptides or fragments thereof can be used as molecular weight  
10 markers. The term "purified" refers to either the "isolated and purified" form of polypeptides of the invention or the "substantially purified" form of polypeptides of the invention, as both are described herein.

A polypeptide "variant" as referred to herein means a polypeptide substantially homologous to native polypeptides of the invention, but which has an amino acid  
15 sequence different from that of native polypeptides (human, murine or other mammalian species) of the invention because of one or more deletions, insertions or substitutions. The variant amino acid sequence preferably is at least 80% identical to a native polypeptide amino acid sequence, most preferably at least 90% identical. The percent identity can be determined, for example, by comparing sequence information using the GAP computer program, version 6.0 described by Devereux et al. (*Nucl. Acids Res.*  
20 12:387, 1984) and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith and Waterman (*Adv. Appl. Math* 2:482, 1981). The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for  
25 nucleotides, and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

25 Variants can comprise conservatively substituted sequences, meaning that a given amino acid residue is replaced by a residue having similar physiochemical characteristics. Examples of conservative substitutions include substitution of one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another, or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other such  
40 conservative substitutions, for example, substitutions of entire regions having similar hydrophobicity characteristics, are well known. Naturally occurring variants are also encompassed by the invention. Examples of such variants are proteins that result from alternate mRNA splicing events, proteolytic cleavage of the polypeptides, or  
45 transcription/translation from different alleles. Variations attributable to proteolysis

5 include, for example, differences in the N- or C-termini upon expression in different types of host cells, due to proteolytic removal of one or more terminal amino acids from the polypeptides (generally from 1-5 terminal amino acids) of the invention.

10 The polypeptides of the invention can also exist as oligomers, such as covalently linked or non-covalently linked dimers or trimers. Oligomers can be linked by disulfide bonds formed between cysteine residues on different polypeptides.

15 In one embodiment of the invention, a polypeptide dimer is created by fusing polypeptides of the invention to the Fc region of an antibody (e.g., IgG1) in a manner that does not interfere with the biological activity of these polypeptides. The Fc region  
20 preferably is fused to the C-terminus of a soluble polypeptide of the invention, to form an Fc fusion or an Fc polypeptide. The terms "Fc fusion protein" or "Fc polypeptides" as used herein includes native and mutein forms, as well as truncated Fc polypeptides containing the hinge region that promotes dimerization. Exemplary methods of making Fc polypeptides set forth above are disclosed in U.S. Patents 5,426,048 and 5,783,672  
25 both of which are incorporated herein by reference.

30 In a preferred embodiment, extracellular domains from transmembrane bound kinases are fused to Fc portions of antibodies to produce soluble Fc polypeptides. These constructs can function as binding sites for the ligand that naturally binds the kinase receptor and thereby inhibit binding of the ligand to the natural receptor.

35 General preparation of fusion proteins comprising heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been described, e.g., by Ashkenazi et al. (*PNAS USA* 88:10535, 1991) and Byrn et al. (*Nature* 344:677, 1990), hereby incorporated by reference. A gene fusion encoding the polypeptide:Fc fusion protein of the invention is inserted into an appropriate expression  
40 vector. Polypeptide:Fc fusion proteins are allowed to assemble much like antibody molecules, whereupon interchain disulfide bonds form between Fc polypeptides, yielding divalent polypeptides of the invention. If fusion proteins are made with both heavy and light chains of an antibody, it is possible to form a polypeptide oligomer with as many as four polypeptides extracellular regions. Alternatively, one can link two soluble  
45 polypeptide domains with a peptide linker.

50 In one embodiment of this invention, the polypeptides of the invention are produced by recombinant expression. In one preferred embodiment, the expression of recombinant polypeptides having kinase functions can be accomplished utilizing fusion of sequences encoding polypeptides having kinase functions to sequences encoding another

5 polypeptide to aid in the purification of polypeptides of the invention. An example of  
such a fusion is a fusion of sequences encoding a polypeptide having kinase functions to  
sequences encoding the product of the *malE* gene of the pMAL-c2 vector of New  
10 England Biolabs, Inc. Such a fusion allows for affinity purification of the fusion protein,  
5 as well as separation of the maltose binding protein portion of the fusion protein from the  
polypeptide of the invention after purification.

15 The insertion of DNA encoding the polypeptide having kinase functions into the  
pMAL-c2 vector can be accomplished in a variety of ways using known molecular  
biology techniques. The preferred construction of the insertion contains a termination  
10 codon adjoining the carboxyl terminal codon of the polypeptide of the invention. In  
addition, the preferred construction of the insertion results in the fusion of the amino  
20 terminus of the polypeptide of the invention directly to the carboxyl terminus of the  
Factor Xa cleavage site in the pMAL-c2 vector. A DNA fragment can be generated by  
PCR using DNA of the invention as the template DNA and two oligonucleotide primers.  
25 Use of the oligonucleotide primers generates a blunt-ended fragment of DNA that can be  
isolated by conventional means. This PCR product can be ligated together with pMAL-  
p2 (digested with the restriction endonuclease *Xmn* I) using conventional means. Positive  
clones can be identified by conventional means. Induction of expression and purification  
30 of the fusion protein can be performed as per the manufacturer's instructions. This  
20 construction facilitates a precise separation of the polypeptide of the invention from the  
fused maltose binding protein utilizing a simple protease treatment as per the  
manufacturer's instructions. In this manner, purified polypeptide having kinase functions  
35 can be obtained. Furthermore, such a constructed vector can be easily modified using  
known molecular biology techniques to generate additional fusion proteins. It is  
25 understood, of course, that many different vectors and techniques can be used for the  
expression and purification of polypeptides of the invention and that this embodiment in  
40 no way limits the scope of the invention.

Polypeptides of the invention can be subjected to fragmentation into peptides by  
chemical and enzymatic means. Chemical fragmentation includes the use of cyanogen  
45 bromide to cleave under neutral or acidic conditions such that specific cleavage occurs at  
methionine residues (E. Gross, *Methods in Enz.* 11:238-255, 1967). This can further  
include additional steps, such as a carboxymethylation step to convert cysteine residues to  
an unreactive species. Enzymatic fragmentation includes the use of a protease such as  
50 Asparaginylendopeptidase, Arginylendo-peptidase, *Achromobacter* protease I, Trypsin,

5 *Staphylococcus aureus* V8 protease, Endoproteinase Asp-N, or Endoproteinase Lys-C under conventional conditions to result in cleavage at specific amino acid residues. Asparaginylendo-peptidase can cleave specifically on the carboxyl side of the asparagine  
10 residues present within the polypeptides of the invention. Arginylendo-peptidase can  
5 cleave specifically on the carboxyl side of the arginine residues present within these polypeptides. *Achromobacter* protease I can cleave specifically on the carboxyl side of the lysine residues present within the polypeptides (Sakiyama and Nakat, U.S. Patent No. 5,248,599; T. Masaki et al., *Biochim. Biophys. Acta* 660:44-50, 1981; T. Masaki et al.,  
15 *Biochim. Biophys. Acta* 660:51-55, 1981). Trypsin can cleave specifically on the  
10 carboxyl side of the arginine and lysine residues present within polypeptides of the invention. *Staphylococcus aureus* V8 protease can cleave specifically on the carboxyl side of the aspartic and glutamic acid residues present within polypeptides (D. W. Cleveland,  
20 *J. Biol. Chem.* 3:1102-1106, 1977). Endoproteinase Asp-N can cleave specifically on the amino side of the asparagine residues present within polypeptides. Endoproteinase Lys-C  
25 can cleave specifically on the carboxyl side of the lysine residues present within polypeptides of the invention. Other enzymatic and chemical treatments can likewise be used to specifically fragment these polypeptides into a unique set of specific peptide molecular weight markers.

30 The resultant fragmented peptides can be analyzed by methods including  
20 sedimentation, electrophoresis, chromatography, and mass spectrometry. The fragmented peptides derived from the polypeptides of the invention can serve as molecular weight markers using such analysis techniques to assist in the determination of the molecular weight of a sample protein. Such a molecular weight determination assists in the  
35 identification of the sample protein. Fragmented peptide molecular weight markers of the  
25 invention are preferably at least 10 amino acids in size. More preferably, these  
40 fragmented peptide molecular weight markers are between 10 and 100 amino acids in size. Even more preferable are fragmented peptide molecular weight markers between 10 and 50 amino acids in size and especially between 10 and 35 amino acids in size. Most preferable are fragmented peptide molecular weight markers between 10 and 20 amino  
45 acids in size.  
30

Furthermore, analysis of the progressive fragmentation of the polypeptides of the invention into specific peptides (D. W. Cleveland et al., *J. Biol. Chem.* 252:1102-1106, 1977), such as by altering the time or temperature of the fragmentation reaction, can be  
50 used as a control for the extent of cleavage of a sample protein. For example, cleavage of

5 the same amount of polypeptide and sample protein under identical conditions can allow  
for a direct comparison of the extent of fragmentation. Conditions that result in the  
complete fragmentation of the polypeptide can also result in complete fragmentation of  
10 the sample protein.

5 In addition, the polypeptides and fragmented peptides of the invention possess  
unique charge characteristics and, therefore, can serve as specific markers to assist in the  
determination of the isoelectric point of a sample protein or fragmented peptide using  
15 techniques such as isoelectric focusing. The technique of isoelectric focusing can be  
further combined with other techniques such as gel electrophoresis to simultaneously  
20 separate a protein on the basis of molecular weight and charge. An example of such a  
combination is that of two-dimensional electrophoresis (T.D. Brock and M.T. Madigan,  
*Biology of Microorganisms* 76-77 (Prentice Hall, 6d ed. 1991)). These polypeptides and  
fragmented peptides thereof can be used in such analyses as markers to assist in the  
determination of both the isoelectric point and molecular weight of a sample protein or  
25 fragmented peptide.

Kits to aid in the determination of apparent molecular weight and isoelectric point  
of a sample protein can be assembled from the polypeptides and peptide fragments of the  
invention. Kits also serve to assess the degree of fragmentation of a sample protein. The  
30 constituents of such kits can be varied, but typically contain the polypeptide and  
fragmented peptide molecular weight markers. Also, such kits can contain the  
polypeptides wherein a site necessary for fragmentation has been removed. Furthermore,  
the kits can contain reagents for the specific cleavage of the polypeptide and the sample  
35 protein by chemical or enzymatic cleavage. Kits can further contain antibodies directed  
against polypeptides or fragments thereof of the invention.

25 The isolated and purified polypeptides of the invention have molecular weights of  
approximately 6883 (HH0900-BF04); 8168 (HH2040-BF04); 39,284 (JJ503-KS); 16,718  
40 (QQ1249-BF04); 58,001 (QQ3351-BF04); 57,381 (SS1771), and 67,331 (SS1771A)  
Daltons in the absence of glycosylation.. The polypeptide of the invention, together with  
a sample protein, can be resolved by denaturing polyacrylamide gel electrophoresis by  
30 conventional means (U. K. Laemmli, *Nature* 227:680-685, 1970) in two separate lanes of  
a gel containing sodium dodecyl sulfate and a concentration of acrylamide between 6-  
20%. Proteins on the gel can be visualized using a conventional staining procedure. The  
polypeptide molecular weight markers of the invention can be used as molecular weight  
50 marker in the estimation of the apparent molecular weight of the sample protein. The

5 unique amino acid sequence of SEQ ID NO:7, 8, 9, 10, 11, 12, and 16 correspond to  
molecular weight of approximately 6883; 8168; 39,284; 16,718; 58,001; 57,381; or  
67,331 Daltons, respectively. Therefore, the polypeptide molecular weight markers serve  
10 particularly well as a molecular weight marker for the estimation of the apparent  
5 molecular weight of sample proteins that have apparent molecular weights close to 6883;  
8168; 39,284; 16,718; 58,001; 57,381; or 67,331 Daltons. The use of these polypeptide  
molecular weight markers allows increased accuracy in the determination of apparent  
15 molecular weight of proteins that have apparent molecular weights close to 6883; 8168;  
39,284; 16,718; 28,982; or 57,381 Daltons. It is understood of course that many different  
10 techniques can be used for the determination of the molecular weight of a sample protein  
using polypeptides of the invention, and that this embodiment in no way limits the scope  
20 of the invention.

Another preferred embodiment of the invention is the use of polypeptides and  
fragmented peptides of the invention as molecular weight markers to estimate the  
25 apparent molecular weight of a sample protein by gel electrophoresis. These fragmented  
15 peptides can be generated methods well known in the art such as chemical fragmentation.  
Isolated and purified polypeptides of the invention can be treated with cyanogen bromide  
under conventional conditions that result in fragmentation of the polypeptide molecular  
30 weight marker by specific hydrolysis on the carboxyl side of the methionine residues  
20 within the polypeptides of the invention (E. Gross, *Methods in Enz.* 11:238-255, 1967).  
Due to the unique amino acid sequence of the polypeptides of the invention, the  
fragmentation of polypeptide molecular weight markers with cyanogen bromide generates  
35 a unique set of fragmented peptide molecular weight markers. The distribution of  
methionine residues determines the number of amino acids in each peptide and the unique  
25 amino acid composition of each peptide determines its molecular weight. Polypeptide  
molecular weight markers of the invention can be analyzed by methods including  
40 sedimentation, gel electrophoresis, chromatography, and mass spectrometry.

The fragmented peptide molecular weight markers of the invention, together with  
a sample protein, can be resolved by denaturing polyacrylamide gel electrophoresis by  
45 conventional means in two separate lanes of a gel containing sodium dodecyl sulfate and  
30 a concentration of acrylamide between 10-20%. Proteins on the gel can be visualized  
using a conventional staining procedure. The fragmented peptide molecular weight  
markers of the invention can be used as molecular weight markers in the estimation of the  
50 apparent molecular weight of the sample protein. The unique amino acid sequence of

5 each marker specifies a molecular weight. Therefore, the fragmented peptide molecular weight markers serve particularly well as molecular weight markers for the estimation of the apparent molecular weight of sample proteins that have similar apparent molecular weights. Consequently, the use of these fragmented peptide molecular weight markers  
10 allows increased accuracy in the determination of apparent molecular weight of proteins.

Polypeptides on the membrane can be visualized using two different methods that allow a discrimination between the sample protein and the molecular weight markers.  
15 Polypeptide or fragmented peptide molecular weight markers of the invention can be visualized using antibodies generated against these markers and conventional immunoblotting techniques. This detection is performed under conventional conditions that do not result in the detection of the sample protein. It is understood that it may not be possible to generate antibodies against all polypeptide fragments of the invention, since small peptides may not contain immunogenic epitopes. It is further understood that not all antibodies will work in this assay; however, those antibodies which are able to bind  
20 polypeptides and fragments of the invention can be readily determined using conventional techniques.

The sample protein is visualized using a conventional staining procedure. The molar excess of sample protein to polypeptide or fragmented peptide molecular weight markers of the invention is such that the conventional staining procedure predominantly  
30 detects the sample protein. The level of these polypeptide or fragmented peptide molecular weight markers is such as to allow little or no detection of these markers by the conventional staining method. The preferred molar excess of sample protein to polypeptide molecular weight markers of the invention is between 2 and 100,000 fold. More preferably, the preferred molar excess of sample protein to these polypeptide  
35 molecular weight markers is between 10 and 10,000 fold and especially between 100 and 1,000 fold.

The polypeptide or fragmented peptide molecular weight markers of the invention can be used as molecular weight and isoelectric point markers in the estimation of the apparent molecular weight and isoelectric point of the sample protein. These polypeptide  
40 or fragmented peptide molecular weight markers serve particularly well as molecular weight and isoelectric point markers for the estimation of apparent molecular weights and isoelectric points of sample proteins that have apparent molecular weights and isoelectric points close to that of the polypeptide or fragmented peptide molecular weight markers of the invention. The ability to simultaneously resolve these polypeptide or fragmented  
45  
50



5 peptide molecular weight markers and the sample protein under identical conditions  
allows for increased accuracy in the determination of the apparent molecular weight and  
isoelectric point of the sample protein. This is of particular interest in techniques, such as  
10 two dimensional electrophoresis, where the nature of the procedure dictates that any  
5 markers should be resolved simultaneously with the sample protein.

In another embodiment, polypeptide or fragmented peptide molecular weight  
15 markers of the invention can be used as molecular weight and isoelectric point markers in  
the estimation of the apparent molecular weight and isoelectric point of fragmented  
peptides derived by treatment of a sample protein with a cleavage agent. It is understood  
20 of course that many techniques can be used for the determination of molecular weight and  
isoelectric point of a sample protein and fragmented peptides thereof using these  
20 polypeptide molecular weight markers and peptide fragments thereof and that this  
embodiment in no way limits the scope of the invention.

Polypeptide molecular weight markers encompassed by invention can have  
25 variable molecular weights, depending upon the host cell in which they are expressed.  
Glycosylation of polypeptide molecular weight markers and peptide fragments of the  
invention in various cell types can result in variations of the molecular weight of these  
markers, depending upon the extent of modification. The size of these polypeptide  
30 molecular weight markers can be most heterogeneous with fragments of polypeptide  
20 derived from the extracellular portion of the polypeptide. Consistent molecular weight  
markers can be obtained by using polypeptides derived entirely from the transmembrane  
and cytoplasmic regions, pretreating with N-glycanase to remove glycosylation, or  
35 expressing the polypeptides in bacterial hosts.

As stated above, the invention provides isolated and purified polypeptides, both  
25 recombinant and non-recombinant. Variants and derivatives of native polypeptides can  
be obtained by mutations of nucleotide sequences coding for native polypeptides.  
40 Alterations of the native amino acid sequence can be accomplished by any of a number of  
conventional methods. Mutations can be introduced at particular loci by synthesizing  
oligonucleotides containing a mutant sequence, flanked by restriction sites enabling  
45 30 ligation to fragments of the native sequence. Following ligation, the resulting  
reconstructed sequence encodes an analog having the desired amino acid insertion,  
substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can  
50 be employed to provide an altered gene wherein predetermined codons can be altered by

substitution, deletion or insertion. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); Kunkel (*Proc. Natl. Acad. Sci. USA* 82:488, 1985); Kunkel et al. (*Methods in Enzymol.* 154:367, 1987); and U.S. Patent Nos. 4,518,584 and 4,737,462, all of which are incorporated by reference.

Polypeptides of the invention can be modified to create polypeptide derivatives by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, polyethylene glycol (PEG) groups, lipids, phosphate, acetyl groups and the like. Covalent derivatives of polypeptides of the invention can be prepared by linking the chemical moieties to functional groups on polypeptide amino acid side chains or at the N-terminus or C-terminus of a polypeptide of the invention or the extracellular domain thereof. Other derivatives of polypeptides within the scope of this invention include covalent or aggregative conjugates of these polypeptides or peptide fragments with other proteins or polypeptides, such as by synthesis in recombinant culture as N-terminal or C-terminal fusions. For example, the conjugate can comprise a signal or leader polypeptide sequence (e.g. the  $\alpha$ -factor leader of *Saccharomyces*) at the N-terminus of a polypeptide of the invention. The signal or leader peptide co-translationally or post-translationally directs transfer of the conjugate from its site of synthesis to a site inside or outside of the cell membrane or cell wall.

Polypeptide conjugates can comprise peptides added to facilitate purification and identification of polypeptides of the invention. Such peptides include, for example, poly-His or the antigenic identification peptides described in U.S. Patent No. 5,011,912 and in Hopp et al., *Bio/Technology* 6:1204, 1988.

The invention further includes polypeptides of the invention with or without associated native-pattern glycosylation. Polypeptides expressed in yeast or mammalian expression systems (e.g., COS-1 or COS-7 cells) can be similar to or significantly different from a native polypeptide in molecular weight and glycosylation pattern, depending upon the choice of expression system. Expression of polypeptides of the invention in bacterial expression systems, such as *E. coli*, provides non-glycosylated molecules. Glycosyl groups can be removed through conventional methods, in particular those utilizing glycopeptidase. In general, glycosylated polypeptides of the invention can be incubated with a molar excess of glycopeptidase (Boehringer Mannheim).

Correspondingly, equivalent DNA constructs that encode various additions or substitutions of amino acid residues or sequences, or deletions of terminal or internal residues or sequences are encompassed by the invention. For example, N-glycosylation sites in the polypeptide extracellular domain can be modified to preclude glycosylation, allowing expression of a reduced carbohydrate analog in mammalian and yeast expression systems. N-glycosylation sites in eukaryotic polypeptides are characterized by an amino acid triplet Asn-X-Y, wherein X is any amino acid except Pro and Y is Ser or Thr. Appropriate substitutions, additions, or deletions to the nucleotide sequence encoding these triplets will result in prevention of attachment of carbohydrate residues at the Asn side chain. Alteration of a single nucleotide, chosen so that Asn is replaced by a different amino acid, for example, is sufficient to inactivate an N-glycosylation site. Known procedures for inactivating N-glycosylation sites in proteins include those described in U.S. Patent 5,071,972 and EP 276,846, hereby incorporated by reference.

In another example, sequences encoding Cys residues that are not essential for biological activity can be altered to cause the Cys residues to be deleted or replaced with other amino acids, preventing formation of incorrect intramolecular disulfide bridges upon renaturation. Other equivalents are prepared by modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present. EP 212,914 discloses the use of site-specific mutagenesis to inactivate KEX2 protease processing sites in a protein. KEX2 protease processing sites are inactivated by deleting, adding, or substituting residues to alter Arg-Arg, Arg-Lys, and Lys-Arg pairs to eliminate the occurrence of these adjacent basic residues. Lys-Lys pairings are considerably less susceptible to KEX2 cleavage, and conversion of Arg-Lys or Lys-Arg to Lys-Lys represents a conservative and preferred approach to inactivating KEX2 sites.

The invention further encompasses isolated fragments and oligonucleotides derived from the nucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 13 or 15. Nucleic acid sequences within the scope of the invention include isolated DNA and RNA sequences that hybridize to the native nucleotide sequences disclosed herein under conditions of moderate or severe stringency, and which encode polypeptides or fragments thereof of the invention. These isolated DNA and RNA sequences also include full length DNA or RNA molecules encoding for polypeptides with kinase activity. As used herein, conditions of moderate stringency, as known to those having ordinary skill in the art, and as defined by Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2 ed.

5 Vol. 1, pp. 1.101-104, Cold Spring Harbor Laboratory Press, (1989), include use of a  
prewashing solution for the nitrocellulose filters 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH  
8.0), hybridization conditions of 50% formamide, 6X SSC at 42°C (or other similar  
10 hybridization solution, such as Stark's solution, in 50% formamide at 42°C), and washing  
5 conditions of about 60°C, 0.5X SSC, 0.1% SDS. Conditions of high stringency are  
defined as hybridization conditions as above, and with washing at 68°C, 0.2X SSC, 0.1%  
SDS. The skilled artisan will recognize that the temperature and wash solution salt  
15 concentration can be adjusted as necessary according to factors such as the length of the  
probe.

10 Due to the known degeneracy of the genetic code, wherein more than one codon  
can encode the same amino acid, a DNA sequence can vary from that shown in SEQ ID  
20 NO: 1, 2, 3, 4, 5, 6, 13 or 15, and still encode a polypeptide having the amino acid  
sequence of SEQ ID NO: 7, 8, 9, 10, 11, 12, 14 or 16. Such variant DNA sequences can  
result from silent mutations (e.g., occurring during PCR amplification), or can be the  
25 product of deliberate mutagenesis of a native sequence.

The invention thus provides equivalent isolated DNA sequences encoding  
polypeptides of the invention, selected from: (a) DNA derived from the coding region of  
a native mammalian gene; (b) cDNA comprising the nucleotide sequence of SEQ ID NO:  
30 1, 2, 3, 4, 5, 6, 13 or 15; (c) DNA encoding the polypeptides of SEQ ID NO: 7, 8, 9, 10,  
20 11, 12, 14 or 16; (d) DNA capable of hybridization to a DNA of (a) under conditions of  
moderate stringency and which encodes polypeptides of the invention; and (e) DNA  
which is degenerate as a result of the genetic code to a DNA defined in (a), (b), (c), or (d)  
35 and which encodes polypeptides of the invention. Of course, polypeptides encoded by  
such DNA equivalent sequences are encompassed by the invention.

25 DNA that is equivalent to the DNA sequence of SEQ ID NO: 1, 2, 3, 4, 5, 6, 13 or  
15 will hybridize under moderately stringent conditions to the double-stranded native  
40 DNA sequence that encode polypeptides comprising amino acid sequences of SEQ ID  
NO: 7, 8, 9, 10, 11, 12, 14 or 16. Examples of polypeptides encoded by such DNA,  
include, but are not limited to, polypeptide fragments and polypeptides comprising  
45 30 inactivated N-glycosylation site(s), inactivated protease processing site(s), or conservative  
amino acid substitution(s), as described above. Polypeptides encoded by DNA derived  
from other mammalian species, wherein the DNA will hybridize to the complement of the  
DNA of SEQ ID NO: 1, 2, 3, 4, 5, 6, 13 or 15 are also encompassed.

5 Recombinant expression vectors containing a nucleic acid sequence encoding polypeptides of the invention can be prepared using well known methods. The expression  
10 vectors include a DNA sequence of the invention operably linked to suitable transcriptional or translational regulatory nucleotide sequences, such as those derived  
5 from a mammalian, microbial, viral, or insect gene. Examples of regulatory sequences include transcriptional promoters, operators, or enhancers, an mRNA ribosomal binding site, and appropriate sequences which control transcription and translation initiation and  
15 termination. Nucleotide sequences are "operably linked" when the regulatory sequence functionally relates to the DNA sequence of the invention. Thus, a promoter nucleotide  
10 sequence is operably linked to a DNA sequence if the promoter nucleotide sequence controls the transcription of the DNA sequence of the invention. The ability to replicate  
20 in the desired host cells, usually conferred by an origin of replication, and a selection gene by which transformants are identified can additionally be incorporated into the expression vector.

15 In addition, sequences encoding appropriate signal peptides that are not naturally associated with polypeptides of the invention can be incorporated into expression vectors. For example, a DNA sequence for a signal peptide (secretory leader) can be fused in-  
30 frame to the nucleotide sequence of the invention so that the polypeptide is initially translated as a fusion protein comprising the signal peptide. A signal peptide that is  
20 functional in the intended host cells enhances extracellular secretion of the polypeptide. The signal peptide can be cleaved from the polypeptide upon secretion of polypeptide from the cell.

35 Suitable host cells for expression of polypeptides of the invention include prokaryotes, yeast or higher eukaryotic cells. Appropriate cloning and expression vectors  
25 for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, in Pouwels et al. *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, (1985). Cell-free translation systems could also be employed to produce polypeptides of  
40 the invention using RNAs derived from DNA constructs disclosed herein.

45 Prokaryotes include gram negative or gram positive organisms, for example, *E. coli* or *Bacilli*. Suitable prokaryotic host cells for transformation include, for example, *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various other species within the  
30 genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. In a prokaryotic host cell, such as *E. coli*, a polypeptide of the invention can include an N-terminal methionine residue to

5 facilitate expression of the recombinant polypeptide in the prokaryotic host cell. The N-terminal Met can be cleaved from the expressed recombinant polypeptide.

10 Expression vectors for use in prokaryotic host cells generally comprise one or more phenotypic selectable marker genes. A phenotypic selectable marker gene is, for example, a gene encoding a protein that confers antibiotic resistance or that supplies an autotrophic requirement. Examples of useful expression vectors for prokaryotic host cells include those derived from commercially available plasmids such as the cloning vector pBR322 (ATCC 37017). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells. To construct 15 an expression vector using pBR322, an appropriate promoter and a DNA sequence of the invention are inserted into the pBR322 vector. Other commercially available vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA). Other commercially available vectors include those that are specifically designed for the expression of proteins; these would 20 include pMAL-p2 and pMAL-c2 vectors that are used for the expression of proteins fused to maltose binding protein (New England Biolabs, Beverly, MA, USA).

Promoter sequences commonly used for recombinant prokaryotic host cell expression vectors include  $\beta$ -lactamase (penicillinase), lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), tryptophan (trp) 30 promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EP-A-36776), and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful prokaryotic host cell expression system employs a phage  $\lambda P_L$  promoter and a cI857ts thermolabile repressor sequence. Plasmid vectors available from the American Type Culture Collection, which incorporate 35 derivatives of the  $\lambda P_L$  promoter, include plasmid pHUB2 (resident in *E. coli* strain JMB9 (ATCC 37092)) and pPLc28 (resident in *E. coli* RR1 (ATCC 53082)).

40 DNA of the invention can be cloned in-frame into the multiple cloning site of an ordinary bacterial expression vector. Ideally the vector contains an inducible promoter upstream of the cloning site, such that addition of an inducer leads to high-level 45 production of the recombinant protein at a time of the investigator's choosing. For some proteins, expression levels can be boosted by incorporation of codons encoding a fusion partner (such as hexahistidine) between the promoter and the gene of interest. The

5 resulting "expression plasmid" can be propagated in a variety of prokaryotic hosts such as *E. coli*.

10 For expression of the recombinant protein, the bacterial cells are propagated in growth medium until reaching a pre-determined optical density. Expression of the  
5 recombinant protein is then induced, e.g. by addition of IPTG (isopropyl-b-D-thiogalactopyranoside), which activates expression of proteins from plasmids containing a lac operator/promoter. After induction (typically for 1-4 hours), the cells are harvested by  
15 pelleting in a centrifuge, e.g. at 5,000 x G for 20 minutes at 4°C.

20 For recovery of the expressed protein, the pelleted cells may be resuspended in ten volumes of 50 mM Tris-HCl (pH 8)/1 M NaCl and then passed two or three times through a French press. Most highly-expressed recombinant proteins form insoluble aggregates known as inclusion bodies. Inclusion bodies can be purified away from the soluble  
25 proteins by pelleting in a centrifuge at 5,000 x G for 20 minutes, 4°C. The inclusion body pellet is washed with 50 mM Tris-HCl (pH 8)/1% Triton X-100 and then dissolved in 50  
15 mM Tris-HCl (pH 8)/8 M urea/ 0.1 M DTT. Any material that cannot be dissolved is removed by centrifugation (10,000 x G for 20 minutes, 20°C). The protein of interest will, in most cases, be the most abundant protein in the resulting clarified supernatant. This protein may be "refolded" into the active conformation by dialysis against 50 mM  
30 Tris-HCl (pH 8)/5 mM CaCl<sub>2</sub>/5 mM Zn(OAc)<sub>2</sub>/1 mM GSSG/0.1 mM GSH. After  
20 refolding, purification can be carried out by a variety of chromatographic methods such as ion exchange or gel filtration. In some protocols, initial purification may be carried out before refolding. As an example, hexahistidine-tagged fusion proteins may be partially  
35 purified on immobilized nickel.

40 While the preceding purification and refolding procedure assumes that the protein  
25 is best recovered from inclusion bodies, those skilled in the art of protein purification will appreciate that many recombinant proteins are best purified out of the soluble fraction of cell lysates. In these cases, refolding is often not required, and purification by standard chromatographic methods can be carried out directly.

45 Polypeptides of the invention alternatively can be expressed in yeast host cells,  
30 preferably from the *Saccharomyces* genus (e.g., *S. cerevisiae*). Other genera of yeast, such as *Pichia*, *K. lactis*, or *Kluyveromyces*, can also be employed. Yeast vectors will often contain an origin of replication sequence from a 2μ yeast plasmid, an autonomously  
50 replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences

5 for transcription termination, and a selectable marker gene. Suitable promoter sequences  
for yeast vectors include, among others, promoters for metallothionein, 3-  
phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980), or other  
10 glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al.,  
5 *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase,  
hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate  
isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase,  
15 phosphoglucose isomerase, and glucokinase. Other suitable vectors and promoters for  
use in yeast expression are further described in Hitzeman, EPA-73,657 or in Fleer et. al.,  
10 *Gene*, 107:285-195 (1991); and van den Berg et. al., *BioTechnology*, 8:135-139 (1990).  
Another alternative is the glucose-repressible ADH2 promoter described by Russell et al.  
20 (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). Shuttle vectors  
replicable in both yeast and *E. coli* can be constructed by inserting DNA sequences from  
pBR322 for selection and replication in *E. coli* (Amp<sup>r</sup> gene and origin of replication) into  
25 the above-described yeast vectors.

The yeast  $\alpha$ -factor leader sequence can be employed to direct secretion of a  
polypeptide of the invention. The  $\alpha$ -factor leader sequence is often inserted between the  
promoter sequence and the structural gene sequence. See, e.g., Kurjan et al., *Cell* 30:933,  
30 1982; Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984; U. S. Patent 4,546,082; and  
20 EP 324,274. Other leader sequences suitable for facilitating secretion of recombinant  
polypeptides from yeast hosts are known to those of skill in the art. A leader sequence  
can be modified near its 3' end to contain one or more restriction sites. This will facilitate  
35 fusion of the leader sequence to the structural gene.

Yeast transformation protocols are known to those of skill in the art. One such  
25 protocol is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929, 1978. The  
Hinnen et al. protocol selects for Trp<sup>r</sup> transformants in a selective medium, wherein the  
40 selective medium consists of 0.67% yeast nitrogen base, 0.5% casamino acids, 2%  
glucose, 10  $\mu$ g/ml adenine, and 20  $\mu$ g/ml uracil.

Yeast host cells transformed by vectors containing ADH2 promoter sequence can  
45 be grown for inducing expression in a "rich" medium. An example of a rich medium is  
one consisting of 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80  
30  $\mu$ g/ml adenine and 80  $\mu$ g/ml uracil. Derepression of the ADH2 promoter occurs when  
glucose is exhausted from the medium.



5 Mammalian or insect host cell culture systems could also be employed to express recombinant polypeptides of the invention. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers,

10 *Bio/Technology* 6:47 (1988). Established cell lines of mammalian origin also can be employed. Examples of suitable mammalian host cell lines include the COS-7 line of monkey kidney cells (ATCC CRL 1651) (Gluzman et al., *Cell* 23:175, 1981), L cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, 15 and BHK (ATCC CRL 10) cell lines, and the CV-1/EBNA-1 cell line (ATCC CRL 10478) derived from the African green monkey kidney cell line CV1 (ATCC CCL 70) as described by McMahan et al. (*EMBO J.* 10: 2821, 1991).

20 Established methods for introducing DNA into mammalian cells have been described (Kaufman, R.J., *Large Scale Mammalian Cell Culture*, 1990, pp. 15-69). Additional protocols using commercially available reagents, such as Lipofectamine (Gibco/BRL) or Lipofectamine-Plus, can be used to transfect cells (Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7417, 1987). In addition, electroporation can be used to transfect mammalian cells using conventional procedures, such as those in Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2 ed. Vol. 1-3, Cold Spring Harbor Laboratory Press, 1989). Selection of stable transformants can be performed using methods known in the art, such as, for example, resistance to cytotoxic drugs. Kaufman et al., *Meth. in Enzymology* 185:487-511, 1990, describes several selection schemes, such as dihydrofolate reductase (DHFR) resistance. A suitable host strain for DHFR selection can be CHO strain DX-B11, which is deficient in DHFR (Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA* 77:4216-4220, 1980). A plasmid expressing the DHFR cDNA can be introduced into strain DX-B11, and only cells that contain the plasmid can grow in the appropriate selective media. Other examples of selectable markers that can be incorporated into an expression vector include cDNAs conferring resistance to antibiotics, such as G418 and hygromycin B. Cells harboring the vector can be selected on the basis of resistance to these compounds.

30 Transcriptional and translational control sequences for mammalian host cell expression vectors can be excised from viral genomes. Commonly used promoter sequences and enhancer sequences are derived from polyoma virus, adenovirus 2, simian virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites can be used to provide other genetic elements for expression of a

5 structural gene sequence in a mammalian host cell. Viral early and late promoters are particularly useful because both are easily obtained from a viral genome as a fragment, which can also contain a viral origin of replication (Fiers et al., *Nature* 273:113, 1978; Kaufman, *Meth. in Enzymology*, 1990).  
10 Smaller or larger SV40 fragments can also be used, provided the approximately 250 bp sequence extending from the *Hind* III site toward the *Bgl* I site located in the SV40 viral origin of replication site is included.

Additional control sequences shown to improve expression of heterologous genes from mammalian expression vectors include such elements as the expression augmenting sequence element (EASE) derived from CHO cells (Morris et al., *Animal Cell*  
15 *Technology*, 1997, pp. 529-534) and the tripartite leader (TPL) and VA gene RNAs from Adenovirus 2 (Gingeras et al., *J. Biol. Chem.* 257:13475-13491, 1982). The internal ribosome entry site (IRES) sequences of viral origin allows dicistronic mRNAs to be translated efficiently (Oh and Sarnow, *Current Opinion in Genetics and Development* 3:295-300, 1993; Ramesh et al., *Nucleic Acids Research* 24:2697-2700, 1996).

20 Expression of a heterologous cDNA as part of a dicistronic mRNA followed by the gene for a selectable marker (e.g. DHFR) has been shown to improve transfectability of the host and expression of the heterologous cDNA (Kaufman, *Meth. in Enzymology*, 1990). Exemplary expression vectors that employ dicistronic mRNAs are pTR-DC/GFP described by Mosser et al., *Biotechniques* 22:150-161, 1997, and p2A5I described by  
25 Morris et al., *Animal Cell Technology*, 1997, pp. 529-534.

A useful high expression vector, pCAVNOT, has been described by Mosley et al., *Cell* 59:335-348, 1989. Other expression vectors for use in mammalian host cells can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983). A useful system for stable high level expression of mammalian cDNAs in C127 murine mammary  
30 epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986). A useful high expression vector, PMLSV N1/N4, described by Cosman et al., *Nature* 312:768, 1984, has been deposited as ATCC 39890. Additional useful mammalian expression vectors are described in EP-A-0367566, and in U.S. Patent Application Serial No. 07/701,415, filed May 16, 1991, incorporated by reference herein.

35 The vectors can be derived from retroviruses. In place of the native signal sequence, a heterologous signal sequence can be added, such as the signal sequence for IL-7 described in United States Patent 4,965,195; the signal sequence for IL-2 receptor described in Cosman et al., *Nature* 312:768 (1984); the IL-4 signal peptide described in  
40 EP 367,566; the type I IL-1 receptor signal peptide described in U.S. Patent 4,968,607;

5 and the type II IL-1 receptor signal peptide described in EP 460,846. Another useful  
expression vector, pFLAG, can be used. FLAG technology is centered on the fusion of a  
low molecular weight (1kD), hydrophilic, FLAG marker peptide to the N-Terminus of a  
10 recombinant protein expressed by the pFLAG-1™ Expression Vector (1) (obtained from  
5 IBI Kodak).

An isolated and purified polypeptide according to the invention can be produced  
by recombinant expression systems as described above or purified from naturally  
15 occurring cells. Polypeptides can be substantially purified, as indicated by a single  
protein band upon analysis by SDS-polyacrylamide gel electrophoresis (SDS-PAGE).

10 One process for producing polypeptides of the invention comprises culturing a  
host cell transformed with an expression vector comprising a DNA sequence that encodes  
20 a polypeptide of the invention under conditions sufficient to promote expression of the  
polypeptide. The polypeptide is then recovered from culture medium or cell extracts,  
depending upon the expression system employed. As is known to the skilled artisan,  
25 procedures for purifying a recombinant protein will vary according to such factors as the  
type of host cells employed and whether or not the recombinant protein is secreted into  
the culture medium. For example, when expression systems that secrete the recombinant  
protein are employed, the culture medium first can be concentrated using a commercially  
30 available protein concentration filter, for example, an Amicon or Millipore Pellicon  
20 ultrafiltration unit. Following the concentration step, the concentrate can be applied to a  
purification matrix such as a gel filtration medium. Alternatively, an anion exchange  
resin can be employed, for example, a matrix or substrate having pendant  
35 diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran,  
cellulose or other types commonly employed in protein purification. Alternatively, a  
25 cation exchange step can be employed. Suitable cation exchangers include various  
insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups  
40 are preferred. Finally, one or more reversed-phase high performance liquid  
chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, (e.g., silica  
gel having pendant methyl or other aliphatic groups) can be employed to further purify  
45 30 the polypeptides. Some or all of the foregoing purification steps, in various  
combinations, are well known and can be employed to provide an isolated and purified  
recombinant protein.

50 It is possible to utilize an affinity column comprising a polypeptide-binding  
protein of the invention, such as a monoclonal antibody generated against polypeptides of

5 the invention, to affinity-purify expressed polypeptides. These polypeptides can be removed from an affinity column using conventional techniques, e.g., in a high salt elution buffer and then dialyzed into a lower salt buffer for use or by changing pH or  
10 other components depending on the affinity matrix utilized.

5 In this aspect of the invention, polypeptide-binding proteins, such as the anti-polypeptide antibodies of the invention, can be bound to a solid phase such as a column chromatography matrix or a similar substrate suitable for identifying, separating, or  
15 purifying cells that express polypeptides of the invention on their surface. Adherence of polypeptide-binding proteins of the invention to a solid phase contacting surface can be accomplished by any means, for example, magnetic microspheres can be coated with  
20 these polypeptide-binding proteins and held in the incubation vessel through a magnetic field. Suspensions of cell mixtures are contacted with the solid phase that has such polypeptide-binding proteins thereon. Cells having polypeptides of the invention on their surface bind to the fixed polypeptide-binding protein and unbound cells then are washed  
25 away. This affinity-binding method is useful for purifying, screening, or separating such polypeptide-expressing cells from solution. Methods of releasing positively selected cells from the solid phase are known in the art and encompass, for example, the use of enzymes. Such enzymes are preferably non-toxic and non-injurious to the cells and are preferably directed to cleaving the cell-surface binding partner.

30 Alternatively, mixtures of cells suspected of containing polypeptide-expressing cells of the invention first can be incubated with a biotinylated polypeptide-binding protein of the invention. Incubation periods are typically at least one hour in duration to ensure sufficient binding to polypeptides of the invention. The resulting mixture then is  
35 passed through a column packed with avidin-coated beads, whereby the high affinity of biotin for avidin provides the binding of the polypeptide-binding cells to the beads. Use of avidin-coated beads is known in the art. See Berenson, et al. *J. Cell. Biochem.*,  
40 10D:239 (1986). Wash of unbound material and the release of the bound cells is performed using conventional methods.

45 In the methods described above, suitable polypeptide-binding proteins are anti-polypeptide antibodies, and other proteins that are capable of high-affinity binding of polypeptides of the invention. A preferred polypeptide-binding protein is an polypeptide monoclonal antibody.

50 Recombinant protein produced in bacterial culture is usually isolated by initial disruption of the host cells, centrifugation, extraction from cell pellets if an insoluble

5 polypeptide, or from the supernatant fluid if a soluble polypeptide, followed by one or more concentration, salting-out, ion exchange, affinity purification or size exclusion chromatography steps. Finally, RP-HPLC can be employed for final purification steps.  
10 Microbial cells can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Transformed yeast host cells are preferably employed to express polypeptides of the invention as a secreted polypeptide in order to simplify purification. Secreted  
15 recombinant polypeptide from a yeast host cell fermentation can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984). Urdal et al.  
20 describe two sequential, reversed-phase HPLC steps for purification of recombinant human IL-2 on a preparative HPLC column.

In yet another embodiment of the invention, antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of  
25 binding to a target mRNA sequence (forming a duplex) or to the sequence in the double-stranded DNA helix (forming a triple helix) can be made according to the invention.  
Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of cDNA (SEQ ID NO: 1, 2, 3, 4, 5, 6, 13 or 15). Such a  
30 fragment generally comprises at least about 14 nucleotides, preferably from about 14 to about 30 nucleotides. The ability to create an antisense or a sense oligonucleotide, based  
20 upon a cDNA sequence for a given protein is described in, for example, Stein and Cohen, *Cancer Res.* 48:2659, 1988 and van der Krol et al., *BioTechniques* 6:958, 1988.

Binding of antisense or sense oligonucleotides to target nucleic acid sequences  
35 results in the formation of complexes that block translation (RNA) or transcription (DNA) by one of several means, including enhanced degradation of the duplexes, premature  
25 termination of transcription or translation, or by other means. The antisense oligonucleotides thus can be used to block expression of polypeptides of the invention.  
40 Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such  
45 oligonucleotides with resistant sugar linkages are stable *in vivo* (i.e., capable of resisting enzymatic degradation), but retain sequence specificity to be able to bind to target nucleotide sequences. Other examples of sense or antisense oligonucleotides include  
50 those oligonucleotides that are covalently linked to organic moieties, such as those described in WO 90/10448, and other moieties that increase affinity of the

oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes can be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

Antisense or sense oligonucleotides can be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example,  $\text{CaPO}_4$ -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. Antisense or sense oligonucleotides are preferably introduced into a cell containing the target nucleic acid sequence by insertion of the antisense or sense oligonucleotide into a suitable retroviral vector, then contacting the cell with the retrovirus vector containing the inserted sequence, either *in vivo* or *ex vivo*. Suitable retroviral vectors include, but are not limited to, the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see PCT Application US 90/02656).

Alternatively, sense or antisense oligonucleotides also can be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

In yet another embodiment, a sense or an antisense oligonucleotide can be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

Another embodiment of the invention relates to therapeutic uses of kinases. Kinases play a central role in cellular signal transduction. As such, alterations in kinase expression and/or activation can have profound effects on a plethora of cellular processes, including, but not limited to, activation or inhibition of cell specific responses, proliferation, and programmed cell death (apoptosis). Over expression of cloned kinases or of catalytically inactive mutants of kinases has been used to identify the role a particular kinase plays in mediating specific signaling events.

5 Kinase mediated cellular signaling often involves a molecular activation cascade,  
during which an activated kinase propagates a ligand-receptor mediated signal by  
specifically phosphorylating target substrates. These substrates can themselves be  
10 kinases which become activated following phosphorylation. Alternatively, they can be  
5 adaptor molecules that facilitate down stream signaling through protein-protein  
interaction following phosphorylation. Regardless of the nature of the substrate  
molecule(s), expressed catalytically active versions of the putative kinases in the  
15 invention can be used to identify what substrate(s) were recognized and phosphorylated  
by the kinase(s) of the invention. As such, these kinases can be used as reagents to  
10 identify novel molecules involved in signal transduction pathways.

20 Knowledge of a particular kinase would enable one to enzymatically label the  
substrate with  $^{32}\text{P}$  thereby facilitating identification and also to use the resultant  
radiolabeled protein or a peptide derived from a certain region of the protein as a  
substrate probe to identify and isolate specific phosphatases. Phosphatases are enzymes  
25 whose function is to remove phosphates from selected phosphoproteins; they perform the  
reverse function of kinases.

30 In some systems specific glycosylations with N-acetyl glucosamine residues have  
been described as a biological counter balance to kinases. That is, glycosidases have been  
suggested to compete with kinases for the same serine or threonine residues to covalently  
20 modify. Therefore, the kinase can be used to dissect dynamic interactions between kinase  
and phosphatase, and also between kinase and glycosidase, and finally to examine the  
dynamic interactions among kinase, phosphatase and glycosidase together.

35 Kinases phosphorylate target serine, threonine or tyrosine residues in the context  
of specific recognition motifs. Recognition motifs can consist solely of primary structure  
or in some cases recognition requires more complex structural features. One can take  
25 advantage of kinases with strict primary sequence recognition requirements by using  
them as a general labeling reagent. Nucleotides coding for the amino acids recognized by  
a particular kinase could be engineered onto either end of a protein on interest, thereby  
40 "tagging" the molecule. The expressed, tagged protein could be  $^{32}\text{P}$ -labeled at a known  
site on the engineered tag by its specific kinase, thus generating a well defined,  
45 radiolabeled protein.

50 Because kinases are phosphotransferases, they must take part in protein-protein  
interactions with at least one or more substrate molecules, i.e. its phosphate recipient(s).  
Therefore, kinases or polypeptides comprised of portions of a kinase could be used as

5 “baits” in the yeast two hybrid system by well established molecular biology techniques, to identify molecules that interact directly with the polypeptide.

Alternatively, polypeptides of the invention could be engineered prior to  
10 expression with a tag such as poly-His or FLAG, then be expressed and purified using either nickel chelate chromatography or anti-FLAG antibody coupled to a resin,  
5 respectively. Once bound to the resin, the polypeptide of the invention could be covalently attached using a bifunctional cross-linking agent using well established  
15 techniques. The covalently bound polypeptide to the resin could then be used to purify molecules from cell lysates or cell supernatants (following treatment with various  
20 reagent) through their affinity for the polypeptide of the invention.

Isolated and purified kinase polypeptides or a fragment thereof of the invention  
20 can also be useful as a therapeutic agent in inhibiting signaling. Polypeptides are introduced into the intracellular environment by well-known means, such as by encasing the protein in liposomes or coupling it to a monoclonal antibody targeted to a specific cell  
25 type.

DNA, polypeptides, and antibodies against polypeptides of the invention can be  
used as reagents in a variety of research protocols. A sample of such research protocols  
are given in Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2 ed. Vol. 1-3,  
30 Cold Spring Harbor Laboratory Press, (1989). For example, these reagents can serve as markers for cell specific or tissue specific expression of RNA or proteins. Similarly,  
20 these reagents can be used to investigate constitutive and transient expression of RNA or polypeptides. The DNA can be used to determine the chromosomal location of DNA and  
35 to map genes in relation to this chromosomal location. The DNA can also be used to examine genetic heterogeneity and heredity through the use of techniques such as genetic  
25 fingerprinting, as well as to identify risks associated with genetic disorders. The DNA can be further used to identify additional genes related to the DNA and to establish  
40 evolutionary trees based on the comparison of sequences. The DNA and polypeptides can be used to select for those genes or proteins that are homologous to the DNA or  
polypeptides, through positive screening procedures such as Southern blotting and  
30 immunoblotting and through negative screening procedures such as subtraction.

45 The polypeptides and fragments of the invention can also be used as a reagent to identify (a) any protein that polypeptide regulates, and (b) other proteins with which it might interact. Polypeptides could be used by coupling recombinant protein to an affinity  
50 matrix, or by using them as a bait in the 2-hybrid system. The polypeptides and



5 fragments thereof can be used as reagents in the study of the kinase signaling pathway as a reagent to block kinase signaling.

10 A hallmark of protein kinases is their ability to phosphorylate other proteins and to auto-phosphorylate. Therefore, in one aspect of the invention, the isolated  
5 polypeptides with kinase activity can be used in assays to phosphorylate target proteins, radiolabel target proteins with  $^{32}\text{P}$ , and identify proteins having phosphatase activity. Exemplary methods of phosphorylation assays set forth above are disclosed in U.S.  
15 Patents 5,447,860 which is incorporated herein by reference. In addition to full length polypeptides, the invention also includes the isolated active kinase domains of kinases,  
10 such as the intracellular domain of transmembrane bound kinases and the cytoplasmic kinases, which can function as reagents in kinase assays. Further, soluble forms of the  
20 extracellular domains of the kinases are useful in inhibiting the natural ligand-receptor interaction.

25 When used as a therapeutic agent, polypeptides of the invention can be formulated  
15 into pharmaceutical compositions according to known methods. The polypeptides can be combined in admixture, either as the sole active material or with other known active materials, with pharmaceutically suitable diluents (e.g., Tris-HCl, acetate, phosphate),  
preservatives (e.g., Thimerosal, benzyl alcohol, parabens), emulsifiers, solubilizers,  
30 adjuvants and/or carriers. Suitable carriers and their formulations are described in  
20 Remington's Pharmaceutical Sciences, 16th ed. 1980, Mack Publishing Co. In addition, such compositions can contain the polypeptides complexed with polyethylene glycol (PEG), metal ions, or incorporated into polymeric compounds such as polyacetic acid,  
35 polyglycolic acid, hydrogels, etc., or incorporated into liposomes, microemulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts or spheroblasts. Such  
25 compositions will influence the physical state, solubility, stability, rate of *in vivo* release, and rate of *in vivo* clearance of polypeptides of the invention.

40 The dosage of the composition can be readily determined by those of ordinary skill in the art. The amount to be administered and the frequency of administration can be determined empirically and will take into consideration the age and size of the patient  
45 30 being treated, as well as the malady being treated.

Treatment comprises administering the composition by any method familiar to those of ordinary skill in the art, including intravenous, intraperitoneal, intracorporeal injection, intra-articular, intraventricular, intrathecal, intramuscular, subcutaneous,  
50 topically, tonsillar, intranasally, intravaginally, and orally. The composition may also be

5 given locally, such as by injection into the particular area, either intramuscularly or subcutaneously.

10 Within the therapeutic and research aspects of the invention, polypeptides of the invention, and peptides based on the amino acid sequence thereof, can be utilized to  
5 prepare antibodies that specifically bind to the polypeptides. The term "antibodies" is meant to include polyclonal antibodies, monoclonal antibodies, fragments thereof such as F(ab')<sub>2</sub>, and Fab fragments, as well as any recombinantly produced binding partners.  
15 Antibodies are defined to be specifically binding if they bind polypeptides of the invention with a  $K_d$  of greater than or equal to about  $10^7$  M<sup>-1</sup>. Affinities of binding  
10 partners or antibodies can be readily determined using conventional techniques, for example those described by Scatchard et al., *Ann. N.Y. Acad. Sci.*, 51:660 (1949).

20 Polyclonal antibodies can be readily generated from a variety of sources, for example, horses, cows, goats, sheep, dogs, chickens, rabbits, mice, or rats, using procedures that are well-known in the art. In general, purified polypeptides of the  
25 invention, or a peptide based on the amino acid sequence of polypeptides of the invention that is appropriately conjugated, is administered to the host animal typically through parenteral injection. The immunogenicity of these polypeptides can be enhanced through the use of an adjuvant, for example, Freund's complete or incomplete adjuvant.

30 Following booster immunizations, small samples of serum are collected and tested for reactivity to the polypeptides. Examples of various assays useful for such determination  
20 include those described in: *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988; as well as procedures such as countercurrent immuno-electrophoresis (CIEP), radioimmunoassay, radio-immunoprecipitation, enzyme-  
35 linked immuno-sorbent assays (ELISA), dot blot assays, and sandwich assays, see U.S. Patent Nos. 4,376,110 and 4,486,530.

40 Monoclonal antibodies can be readily prepared using well-known procedures, see for example, the procedures described in U.S. Patent Nos. RE 32,011, 4,902,614, 4,543,439, and 4,411,993; *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Plenum Press, Kennett, McKearn, and Bechtol (eds.), 1980. Briefly,  
45 the host animals, such as Balb/c mice are injected intraperitoneally at least once, and preferably at least twice at about 3 week intervals with isolated and purified polypeptides or conjugated polypeptides of the invention, optionally in the presence of adjuvant. 10  
50  $\mu$ g of isolated and purified polypeptide of the invention or peptides based on the amino acid sequence of polypeptides of the invention in the presence of RIBI adjuvant (RIBI

5 Corp., Hamilton, Montana). Mouse sera are then assayed by conventional dot blot  
technique or antibody capture (ABC) to determine which animal produces the highest  
level of antibody and whose spleen cells are the best candidate for fusion. Approximately  
10 two to three weeks later, the mice are given an intravenous boost of the polypeptides or  
5 conjugated polypeptides such as 3µg suspended in sterile PBS. Mice are later sacrificed  
and spleen cells fused with commercially available myeloma cells, such as Ag8.653  
(ATCC), following established protocols. Briefly, the myeloma cells are washed several  
15 times in media and fused to mouse spleen cells at a ratio of about three spleen cells to one  
myeloma cell. The fusing agent can be any suitable agent used in the art, for example,  
10 polyethylene glycol (PEG) or more preferably, 50% PEG: 10% DMSO (Sigma). Fusion  
is plated out into, for example, twenty 96-well flat bottom plates (Corning) containing an  
appropriate medium, such as HAT supplemented DMEM media and allowed to grow for  
20 eight days. Supernatants from resultant hybridomas are collected and added to, for  
example, a 96-well plate for 60 minutes that is first coated with goat anti-mouse Ig.  
25 Following washes, <sup>125</sup>I-polypeptide or peptides of the invention are added to each well,  
incubated for 60 minutes at room temperature, and washed four times. Positive wells can  
be subsequently detected by conventional methods, such as autoradiography at -70°C  
using Kodak X-Omat S film. Positive clones can be grown in bulk culture and  
30 supernatants are subsequently purified, such as over a Protein A column (Pharmacia). It  
20 is understood of course that many techniques could be used to generate antibodies against  
polypeptides and fragmented peptides of the invention and that this embodiment in no  
way limits the scope of the invention.

35 The monoclonal antibodies of the invention can be produced using alternative  
techniques, such as those described by Altling-Mees et al., "Monoclonal Antibody  
25 Expression Libraries: A Rapid Alternative to Hybridomas", *Strategies in Molecular  
Biology* 3:1-9 (1990), which is incorporated herein by reference. Similarly, binding  
40 partners can be constructed using recombinant DNA techniques to incorporate the  
variable regions of a gene that encodes a specific binding antibody. Such a technique is  
described in Larrick et al., *Biotechnology*, 7:394 (1989).

45 30 Other types of "antibodies" can be produced using the information provided herein  
in conjunction with the state of knowledge in the art. For example, antibodies that have  
been engineered to contain elements of human antibodies that are capable of specifically  
binding polypeptides of the invention are also encompassed by the invention.

5                   Once isolated and purified, the antibodies against polypeptides of the invention  
can be used to detect the presence of the polypeptides in a sample using established assay  
protocols. Further, the antibodies of the invention can be used therapeutically or for  
10 research purposes to bind to the polypeptides and inhibit its activity *in vivo* or *in vitro*.

5                   Antibodies immunoreactive with polypeptides of the invention, and in particular,  
monoclonal antibodies against these polypeptides, are now made available through the  
invention. Such antibodies can be useful for inhibiting polypeptide activity *in vivo* and  
15 for detecting the presence of polypeptides of the invention in a sample.

                  In another embodiment, antibodies generated against a polypeptide and  
10 fragmented peptides of the invention can be used in combination with polypeptide or  
fragmented peptide molecular weight markers of the invention to enhance the accuracy in  
20 the use of these molecular weight markers to determine the apparent molecular weight  
and isoelectric point of a sample protein. Polypeptide or fragmented peptide molecular  
weight markers of the invention can be mixed with a molar excess of a sample protein  
25 and the mixture can be resolved by two dimensional electrophoresis by conventional  
means. Polypeptides can be transferred to a suitable protein binding membrane, such as  
nitrocellulose, by conventional means and detected by the antibodies of the invention.

                  The purified polypeptides according to the invention will facilitate the discovery  
30 of inhibitors of such polypeptides. The use of a purified polypeptide of the invention in  
20 the screening of potential inhibitors thereof is important and can eliminate or reduce the  
possibility of interfering reactions with contaminants.

                  In addition, polypeptides of the invention can be used for structure-based design  
35 of polypeptide-inhibitors. Such structure-based design is also known as "rational drug  
design." The polypeptides can be three-dimensionally analyzed by, for example, X-ray  
25 crystallography, nuclear magnetic resonance or homology modeling, all of which are  
well-known methods. The use of the polypeptide structural information in molecular  
40 modeling software systems to assist in inhibitor design and inhibitor- polypeptide  
interaction is also encompassed by the invention. Such computer-assisted modeling and  
drug design can utilize information such as chemical conformational analysis,  
45 30 electrostatic potential of the molecules, protein folding, etc. For example, most of the  
design of class-specific inhibitors of metalloproteases has focused on attempts to chelate  
or bind the catalytic zinc atom. Synthetic inhibitors are usually designed to contain a  
negatively-charged moiety to which is attached a series of other groups designed to fit the  
50 specificity pockets of the particular protease. A particular method of the invention

5 comprises analyzing the three dimensional structure of polypeptides of the invention for likely binding sites of substrates, synthesizing a new molecule that incorporates a predictive reactive site, and assaying the new molecule as described above.

10 The polypeptides of the present invention may also be used in a screening assay to identify compounds and small molecules which inhibit (antagonize) or enhance (agonize) activation of the polypeptides of the instant invention. Thus, for example, polypeptides of the invention may be used to identify antagonists and agonists from cells, cell-free preparations, chemical libraries, and natural product mixtures. The antagonists and agonists may be natural or modified substrates, ligands, enzymes, receptors, etc. of the polypeptides of the instant invention, or may be structural or functional mimetics of the polypeptides. Potential antagonists of the polypeptides of the instant invention may include small molecules, peptides, and antibodies that bind to and occupy a binding site of the polypeptides, causing them to be unavailable to bind to their ligands and therefore preventing normal biological activity. Other potential antagonists are antisense molecules which may hybridize to mRNA *in vivo* and block translation of the mRNA into the polypeptides of the instant invention. Potential agonists include small molecules, peptides and antibodies which bind to the instant polypeptides and elicit the same or enhanced biological effects as those caused by the binding of the polypeptides of the instant invention.

20 Small molecule agonists and antagonists are usually less than 10K molecular weight and may possess a number of physiochemical and pharmacological properties that enhance cell penetration, resist degradation and prolong their physiological half-lives. (Gibbs, J., *Pharmaceutical Research in Molecular Oncology, Cell*, Vol. 79 (1994).) Antibodies, which include intact molecules as well as fragments such as Fab and F(ab')<sub>2</sub> fragments, may be used to bind to and inhibit the polypeptides of the instant invention by blocking the commencement of a signaling cascade. It is preferable that the antibodies are humanized, and more preferable that the antibodies are human. The antibodies of the present invention may be prepared by any of a variety of well-known methods.

30 Specific screening methods are known in the art and many are extensively incorporated in high throughput test systems so that large numbers of test compounds can be screened within a short amount of time. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, cell based assays, etc. These assay formats are well known in the art. The screening assays of the present invention are amenable to screening of chemical

5 libraries and are suitable for the identification of small molecule drug candidates, antibodies, peptides and other antagonists and agonists.

10 One embodiment of a method for identifying molecules which antagonize or inhibit the polypeptides involves adding a candidate molecule to a medium which  
5 contains cells that express the polypeptides of the instant invention; changing the conditions of said medium so that, but for the presence of the candidate molecule, the polypeptides would be bound to their ligands; and observing the binding and stimulation  
15 or inhibition of a functional response. The activity of the cells which were contacted with the candidate molecule may then be compared with the identical cells which were not  
10 contacted and agonists and antagonists of the polypeptides of the instant invention may be identified. The measurement of biological activity may be performed by a number of  
20 well-known methods such as measuring the amount of protein present (e.g. an ELISA) or of the protein's activity. A decrease in biological stimulation or activation would indicate an antagonist. An increase would indicate an agonist. Specifically, one  
25 embodiment of the instant invention includes agonists and antagonists of QQ3351, SS1771, SS1771A and truncated QQ3351.

Screening assays can further be designed to find molecules that mimic the  
30 biological activity of the polypeptides of the instant invention. Molecules which mimic the biological activity of a polypeptide may be useful for enhancing the biological activity  
20 of the polypeptide. To identify compounds for therapeutically active agents that mimic the biological activity of a polypeptide, it must first be determined whether a candidate molecule binds to the polypeptide. A binding candidate molecule is then added to a  
35 biological assay to determine its biological effects. The biological effects of the candidate molecule are then compared to the those of the polypeptide.

25 The specification is most thoroughly understood in light of the teachings of the references cited within the specification, which are hereby incorporated by reference.  
40 The embodiments within the specification provide an illustration of embodiments of the invention and should not be construed to limit the scope of the invention. The skilled artisan recognizes many other embodiments are encompassed by the claimed invention.

## Claims

5

10

15

20

25

30

35

40

45

50

55

5 What is claimed is:

1. An isolated nucleic acid molecule comprising the DNA of SEQ ID NO:1, 2, 3, 4, 5, 6, 13 and 15.

10 2. An isolated nucleic acid molecule encoding an amino acid sequence comprising the sequence of SEQ ID NO:7, 8, 9, 10, 11, 12, 14 and 16.

15 3. An isolated nucleic acid molecule that hybridizes to either strand of a denatured, double-stranded DNA comprising the nucleic acid sequence of any one of claims 1 or 2 under conditions of moderate stringency in 50% formamide and 6XSSC, at 42°C with washing conditions of 60°C, 0.5XSSC, 0.1% SDS.

20 4. The isolated nucleic acid molecule as claimed in claim 3, wherein said isolated nucleic acid molecule is derived by *in vitro* mutagenesis from SEQ ID NO: 1, 2, 3, 4, 5, 6, 13 and 15.

25 5. An isolated nucleic acid molecule degenerate from SEQ ID NO: 1, 2, 3, 4, 5, 6, 13 and 15 as a result of the genetic code.

30 6. An isolated nucleic acid molecule, which is human DNA, an allelic variant of human DNA, or a species homolog of DNA selected from the group consisting of HH0900, HH2040, JJ503, QQ1249, QQ3351, QQ3351 (truncated), SS1771, and SS1771A.

35 7. A recombinant vector that directs the expression of a nucleic acid molecule selected from the group consisting of the nucleic acid molecules of claims 1, 2, 5, and 6.

40 8. A recombinant vector that directs the expression of a nucleic acid molecule of claim 3.

45 9. A recombinant vector that directs the expression of a nucleic acid molecule of claim 4.



5                   10.    An isolated polypeptide encoded by a nucleic acid molecule selected from the group consisting of the nucleic acid molecules of claims 1, 2, 5, and 6.

10                  11.    An isolated polypeptide according to claim 10 having a molecular weight of approximately 6883; 8168; 39,284; 16,718; 58,001; 57,381; and 67,331 Daltons as determined by SDS-PAGE.

15                  12.    An isolated polypeptide according to claim 10 in non-glycosylated form.

                  13.    An isolated polypeptide encoded by a nucleic acid molecule of claim 3.

20                  14.    An isolated polypeptide according to claim 13 in non-glycosylated form.

                  15.    An isolated polypeptide encoded by a nucleic acid molecule of claim 4.

25                  16.    An isolated polypeptide according to claim 15 in non-glycosylated form.

                  17.    Isolated antibodies that bind to a polypeptide of claim 10.

30                  18.    Isolated antibodies according to claim 17, wherein the antibodies are monoclonal antibodies.

35                  19.    Isolated antibodies that bind to a polypeptide of claim 13.

                  20.    Isolated antibodies according to claim 19, wherein the antibodies are monoclonal antibodies.

40                  21.    Isolated antibodies that bind to a polypeptide of claim 15.

45                  22.    Isolated antibodies according to claim 21, wherein the antibodies are monoclonal antibodies.

                  23.    A host cell transfected or transduced with the vector of claim 7.

5                   24.     A method for the production of polypeptide comprising culturing a host cell of claim 23 under conditions promoting expression, and recovering the polypeptide from the culture medium.

10                   25.     The method of claim 24, wherein the host cell is selected from the group consisting of bacterial cells, yeast cells, plant cells, insect cells, and animal cells.

15                   26.     A host cell transfected or transduced with the vector of claim 8.

20                   27.     A method for the production of polypeptide comprising culturing a host cell of claim 26 under conditions promoting expression, and recovering the polypeptide from the culture medium.

25                   28.     The method of claim 27, wherein the host cell is selected from the group consisting of bacterial cells, yeast cells, plant cells, insect cells, and animal cells.

                    29.     A host cell transfected or transduced with the vector of claim 9.

30                   30.     A method for the production of polypeptide comprising culturing a host cell of claim 29 under conditions promoting expression, and recovering the polypeptide from the culture medium.

35                   31.     The method of claim 30, wherein the host cell is selected from the group consisting of bacterial cells, yeast cells, plant cells, and animal cells.

40                   32.     A method of screening a candidate molecule to identify its ability to inhibit (antagonize) or agonize a polypeptide encoded by cDNA selected from the group consisting of SEQ. ID. NO.:1, 2, 3, 4, 5, 6, 13 and 15, said method comprising the steps of:

- 45                   (a)     adding said candidate molecule to a medium which contains cells expressing said polypeptide;
- (b)     changing the conditions of said medium so that, but for the presence of said candidate molecule, said polypeptide would be bound;

- 5
- (c) determining the level of biological activity of said polypeptide formed in said medium; and
- (d) comparing the level of biological activity of step (c) with the level of biological activity that occurs in said medium in the absence of said candidate molecule;
- 10

wherein a decreased level of biological activity of step (c) as compared to the level of biological activity that occurs in said medium in the absence of said candidate molecule indicates an antagonist and an increased level of biological activity indicates an agonist.

15

33. An antagonist as identified by the method of claim 32.

20

34. The antagonist of claim 33, wherein said cDNA is selected from the group consisting of SEQ.ID. NO.: 5, 6, 13 and 15.

25

35. An agonist as identified by the method of claim 32.

36. A method of screening a candidate molecule to identify its ability to mimic the biological activity of a polypeptide encoded by cDNA selected from the group consisting of SEQ. ID. NO.:1, 2, 3, 4, 5, 6, 13 or 15, said method comprising the steps of :

30

a. determining if said candidate molecule binds to said polypeptide or fragments thereof;

35

b. adding said candidate molecule to a biological assay to determine its biological effects; and

c. comparing said biological effects of said candidate molecule with the biological effects of said polypeptide.

40

## SEQUENCE LISTING

<110> Virca, Duke  
 Bird, Timothy A.  
 5 Anderson, Dirk M.  
 Marken, John S.  
 Immunex Corporation

<120> Human DNA and Polypeptides Having Kinase Functions

10 <130> 2877-WO

<140> --to be assigned--  
 <141> 1999-08-03

15 <150> 60/099,972  
 <151> 1998-09-11

<150> 60/095,270  
 20 <151> 1998-08-04

<160> 12

<170> PatentIn Ver. 2.0

25 <210> 1  
 <211> 181  
 <212> DNA  
 <213> Homo sapiens

30 <400> 1  
 gtacgccatg aaggtgctgc gcaaggcggc gctggtgcag cgcgcccaaga cgcaagagca 60  
 cagcgccacc gagcgctcgg tgctggagct ggtgcgccag gcgcccttcc tggtcacgct 120  
 gcactacgct ttccagacgg atgccaagct gcacctcatc ctggactatg tgagcggcgg 180  
 35 g 181

<210> 2  
 <211> 221  
 <212> DNA  
 40 <213> Homo sapiens

<400> 2  
 cccgagaggt gccacatcag accgcctccg acttcgtgcg ggactcggcg gccagccacc 60  
 aggcggagcc cgagggcgtag gagcggcgcg tgtgcttccg gcttctgcaa ctctgcaacg 120  
 45 ggctggagca cctgaaggag caggggatca tccaccggga cctgtgcctg gagaacctgc 180  
 tgctggtgca ctgcacctc caggccggcc cggggcccgc c 221

<210> 3  
 <211> 1085  
 50 <212> DNA  
 <213> Homo sapiens

<400> 3  
 cgggcagggc tggagctggg ctgggatccc gagctcggca gcagcgcagc gggccggccc 60  
 55 acctgctggt gccttgagg ctctgagccc cggcggcgcc cgggcccacg cggaacgacg 120  
 gggcgagatg cgagccaccc ctctggctgc tcctgcgggt tcctgtcca ggaagaagcg 180  
 gttggagttg gatgacaact tagataccga gcgtcccgtc cagaaacgag ctccaagtgg 240  
 gccccagccc agactgcccc cctgcctgtt gcccctgagc ccacctactg ctccagatcg 300  
 tgcaactgct gtggccactg cctcccgtct tgggccctat gtctcctctg agcccaggga 360  
 60 gggcgggccc gcctaccagg ccctgcactg ccctacaggc actgagtata cctgcaaggt 420

```

gtaccccgctc caggaagccc tggccgtgct ggagccctac gcgcggctgc ccccgacaaa 480
gcatgtggctt cggccactg aggtcctggc tggtaaccag ctccctctacg cctttttcac 540
tcggaccat ggggacatgc acagcctggg gcgaagccgc caccgtatcc ctgagcctga 600
ggctgccgtg ctcttccgcc agatggccac cgccctggcg cactgtcacc agcacggtct 660
ggctctcgct gatctcaagc tgtgtcgctt tgtcttcgct gaccgtgaga ggaagaagct 720
gggtgctggag aacctggagg actcctgcgt gctgactggg ccagatgatt ccctgtggga 780
caagcacgcg tgcccagcct acgtgggacc tgagatactc agctcacggg cctcactact 840
gggcaaggca gccgatgtct ggagcctggg cgtggcgctc ttcacatgc tggccggcca 900
ctaccccttc caggactcgg agcctgtcct gctcttcggc aagatccgcc gcggggccta 960
cgcccttgct gcaggcctct cggccctgc cgcctgtctg gtctgctgcc tccttcgtcg 1020
ggagccagct gaacggctca cagccacagg catcctcctg caccctggc tcgcacagga 1080
cccga 1085

```

&lt;210&gt; 4

&lt;211&gt; 388

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

```

cagcgagaag ccgacatgca tcgcctcttc aatcacccca acatccttcg cctcgtggct 60
tactgtctga gggaaacggg tgctaagcat gaggcctggc tgctgctacc attcttcaag 120
agaggtacgc tgtggaatga gatagaaagg ctgaaggaca aaggcaactt cctgaccgag 180
gatcaaatcc tttggctgct gctggggatc tgcagaggcc ttgaggccat tcatgccaag 240
ggttatgcct acagagactt gaagcccacc aatatattgc ttggagatga ggggcagcca 300
gttttaatgg acttgggttc catgaatcaa gcatgcatcc atgtggaggg ctcccggcag 360
gctctgacct tgcaggactg ggcagccc 388

```

&lt;210&gt; 5

&lt;211&gt; 1555

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

```

atgctaacta gtttaaacag atcttgaac gagacgacct gctgtggaag agcgagcttt 60
ttggaactgt gcacgggaca gattggacgc acaccctcgc ggagcgcgga aggcattgaa 120
aatttgaagc atattatcac ccttggccag gtcateccaca aacgggtgtga agagatgaaa 180
tactgcaaga aacagtgcgc gcgcctgggc caccgcgtcc tcggcctgat caagcctctg 240
gagatgctcc aggaccaagg aaagaggagc gtgcctctcg agaagttaac cacagccatg 300
aaccgcttca aggctgccct ggaggaggct aatggggaga tagaaaagt cagcaataga 360
tccaatatct gcaggtttct aacagcaagc caggacaaaa tactcttcaa ggacgtgaac 420
aggaagctga gtgatgtctg gaaggagctc tcgctgttac ttcaggttga gcaacgcattg 480
cctgtttcac ccataagcca aggagcgtcc tgggcacagg aagatcagca ggatgcagac 540
gaagacaggc gagctttcca gatgctaaga agagataatg aaaaaataga agcttactg 600
agacgattag aaatcaacat gaaagaaatc aaggaaactt tgaggcagta ttaccacca 660
aaatgcattg aggagatccc gcaagagcaa atcaaggaga tcaagaagga gcagctttca 720
ggatccccgt ggattctgct aagggaaaat gaagtcagca cactttataa aggagaatac 780
cacagagctc cagtggccat aaaagtattc aaaaaactcc aggcctggcag cattgcaata 840
gtgaggcaga ctttcaataa ggagatcaaa accatgaaga aattcgaatc tcccaacatc 900
ctgcgtatat ttgggatttg cattgatgaa acagtgactc cgcctcaatt ctccattgtc 960
atggagtact gtgaactcgg gaccctgagg gagctgttgg atagggaaaa agacctcaca 1020
cttggcaagc gcatgtcct agtcctgggg gcagcccgag gcctataccg gctacaccat 1080
tcagaagcac ctgaactcca cggaaaaatc agaagctcaa acttcctggt aactcaaggc 1140
taccaagtga agcttgcagg atttgagttg aggaaaacac agacttccat gaggttggga 1200
actacgagag aaaagacaga cagagtcaaa tctacagcat atctctcacc tcaggaactg 1260
gaagatgtat ttatcaata tgatgtaaag tctgaaatat acagcttttg aatcgtcctc 1320
tggaataatc ccactggaga tatcccgttt caaggctgta attctgagaa gatccgcaag 1380
ctggtggctg tgaagcgcca gcaggagcca ctgggtgaag actgccttc agagctcgg 1440
gagatcattg atgagtgcgc ggcagcaggt cgtctcgttc caagatctgt agcggccgac 1500
cgggcgctgc acgttttaac gcgtggccct cgagaggttt tccgatccgg tcgat 1555

```

<210> 6  
 <211> 1498  
 <212> DNA  
 <213> Homo sapiens

<400> 6  
 cttcccgcgtg gacgtggagt acggaggccc agaccggagg tgcccgcctc cgccttacc 60  
 gaagcacctg ctgctgcgca gcaagtcgga gcagtagcac ctggacagcc tgtgcgcagg 120  
 catggagcag agcctccgtg cgggcccctaa cgagcccagag ggcggcgaca agagccgcaa 180  
 aagcgccaag ggggacaaag gcggaaagga taaaaagcag attcagacct ctcccgttcc 240  
 cgtccgcaaa aacagcagag acgaagagaa gagagagtca cgcataaaga gctactgtcc 300  
 atacgccttt aagttcttca tggagcagca cgtggagaat gtcataaaaa cctaccagca 360  
 gaagggttaac cggaggctgc agctggagca agaaatggcc aaagctggac tctgtgaagc 420  
 tgagcaggag cagatgcgga agatcctcta ccagaaagag tctaattaca acagggttaa 480  
 gagggccaag atggacaagt ctatgtttgt caagatcaaa accctgggga tcggtgcctt 540  
 tggagaagtg tgccttgctt gtaagggtga cactcacgcc ctgtacgcca tgaagaccct 600  
 aaggaaaaag gatgtcctga accggaatca ggtggccac gtcaaggccg agagggacat 660  
 cctggccgag gcagacaatg agtgggtggt caaactctac tactccttcc aagacaaaga 720  
 cagcctgtac tttgtgatgg actacatccc tgggtgggac atgatgagcc tgcgtatccc 780  
 gatggaggtc ttcccgtgag acctggcccg gttctacatc gcagagctga ctttggccat 840  
 tgagagtgtc cacaagatgg gttcatcca ccgagacatc aagcctgata acattttgat 900  
 agatctggat ggtcacatta aactcacaga ttctggcctc tgcactgggt tcagggtggac 960  
 tcacaattcc aatatattacc agaaggagg ccattgtcaga caggacagca tggagcccag 1020  
 cgacctctgg gatgatgtgt ctaactgtcg gtgtggggac aggtgaaga ccctagagca 1080  
 gagggcgcgg aagcagcacc agaggtgcct ggcacattca ctgggtggga ctccaaacta 1140  
 catcgacccc gaggtgtctc tccgcaaagg gtacactcaa ctctgtgact ggtggagtgt 1200  
 tggagtgatt ctcttcgaga tgctgggtgg gcagccgccc tttttggcac ctactccac 1260  
 agaaaccag ctgaaggatg tcaactggga gaacacgctc cacattccag cccaggtgaa 1320  
 gctgagccct gagggcaggg acctcatcac caagctgtgc tgctccgcag accaccgctt 1380  
 gggggcgaat gggggcagat acctgaaggc ccacccttc ttcagcgcca ttgacttctc 1440  
 cagtgcacac cgaagcacc cagccccta cgttcccacc atcagccacc ccatggag 1498

<210> 7  
 <211> 60  
 <212> PRT  
 <213> Homo sapiens

<400> 7  
 Tyr Ala Met Lys Val Leu Arg Lys Ala Ala Leu Val Gln Arg Ala Lys  
 1 5 10 15  
 Thr Gln Glu His Thr Arg Thr Glu Arg Ser Val Leu Glu Leu Val Arg  
 20 25 30  
 Gln Ala Pro Phe Leu Val Thr Leu His Tyr Ala Phe Gln Thr Asp Ala  
 35 40 45  
 Lys Leu His Leu Ile Leu Asp Tyr Val Ser Gly Gly  
 50 55 60

<210> 8  
 <211> 73  
 <212> PRT  
 <213> Homo sapiens

<400> 8  
 Arg Glu Val Pro His Gln Thr Ala Ser Asp Phe Val Arg Asp Ser Ala  
 1 5 10 15

Ala Ser His Gln Ala Glu Pro Glu Ala Tyr Glu Arg Arg Val Cys Phe  
                   20                  25                  30  
 Leu Leu Leu Gln Leu Cys Asn Gly Leu Glu His Leu Lys Glu His Gly  
                   35                  40                  45  
 Ile Ile His Arg Asp Leu Cys Leu Glu Asn Leu Leu Leu Val His Cys  
                   50                  55                  60  
 Thr Leu Gln Ala Gly Pro Gly Pro Ala  
                   65                  70

<210> 9  
 <211> 360  
 <212> PRT  
 <213> Homo sapiens

<400> 9  
 Gly Gln Gly Trp Ser Trp Ala Gly Ile Pro Ser Ser Ala Ala Ala Gln  
                   1                  5                  10                  15  
 Arg Ala Gly Pro Pro Ala Gly Ala Leu Glu Ala Leu Ser Pro Gly Gly  
                   20                  25                  30  
 Ala Arg Ala His Ala Glu Arg Arg Gly Glu Met Arg Ala Thr Pro Leu  
                   35                  40                  45  
 Ala Ala Pro Ala Gly Ser Leu Ser Arg Lys Lys Arg Leu Glu Leu Asp  
                   50                  55                  60  
 Asp Asn Leu Asp Thr Glu Arg Pro Val Gln Lys Arg Ala Arg Ser Gly  
                   65                  70                  75                  80  
 Pro Gln Pro Arg Leu Pro Pro Cys Leu Leu Pro Leu Ser Pro Pro Thr  
                   85                  90                  95  
 Ala Pro Asp Arg Ala Thr Ala Val Ala Thr Ala Ser Arg Leu Gly Pro  
                   100                  105                  110  
 Tyr Val Leu Leu Glu Pro Glu Glu Gly Gly Arg Ala Tyr Gln Ala Leu  
                   115                  120                  125  
 His Cys Pro Thr Gly Thr Glu Tyr Thr Cys Lys Val Tyr Pro Val Gln  
                   130                  135                  140  
 Glu Ala Leu Ala Val Leu Glu Pro Tyr Ala Arg Leu Pro Pro His Lys  
                   145                  150                  155                  160  
 His Val Ala Arg Pro Thr Glu Val Leu Ala Gly Thr Gln Leu Leu Tyr  
                   165                  170                  175  
 Ala Phe Phe Thr Arg Thr His Gly Asp Met His Ser Leu Val Arg Ser  
                   180                  185                  190  
 Arg His Arg Ile Pro Glu Pro Glu Ala Ala Val Leu Phe Arg Gln Met  
                   195                  200                  205  
 Ala Thr Ala Leu Ala His Cys His Gln His Gly Leu Val Leu Arg Asp  
                   210                  215                  220

Leu Lys Leu Cys Arg Phe Val Phe Ala Asp Arg Glu Arg Lys Lys Leu  
 225 230 235 240  
 Val Leu Glu Asn Leu Glu Asp Ser Cys Val Leu Thr Gly Pro Asp Asp  
 245 250 255  
 Ser Leu Trp Asp Lys His Ala Cys Pro Ala Tyr Val Gly Pro Glu Ile  
 260 265 270  
 Leu Ser Ser Arg Ala Ser Tyr Ser Gly Lys Ala Ala Asp Val Trp Ser  
 275 280 285  
 Leu Gly Val Ala Leu Phe Thr Met Leu Ala Gly His Tyr Pro Phe Gln  
 290 295 300  
 Asp Ser Glu Pro Val Leu Leu Phe Gly Lys Ile Arg Arg Gly Ala Tyr  
 305 310 315 320  
 Ala Leu Pro Ala Gly Leu Ser Ala Pro Ala Arg Cys Leu Val Arg Cys  
 325 330 335  
 Leu Leu Arg Arg Glu Pro Ala Glu Arg Leu Thr Ala Thr Gly Ile Leu  
 340 345 350  
 Leu His Pro Trp Leu Arg Gln Asp  
 355 360

<210> 10  
 <211> 146  
 <212> PRT  
 <213> Homo sapiens

<400> 10  
 Gln Arg Glu Ala Asp Met His Arg Leu Phe Asn His Pro Asn Ile Leu  
 1 5 10 15  
 Arg Leu Val Ala Tyr Cys Leu Arg Glu Arg Gly Ala Lys His Glu Ala  
 20 25 30  
 Trp Leu Leu Leu Pro Phe Phe Lys Arg Gly Thr Leu Trp Asn Glu Ile  
 35 40 45  
 Glu Arg Leu Lys Asp Lys Gly Asn Phe Leu Thr Glu Asp Gln Ile Leu  
 50 55 60  
 Trp Leu Leu Leu Gly Ile Cys Arg Gly Leu Glu Ala Ile His Ala Lys  
 65 70 75 80  
 Gly Tyr Ala Tyr Arg Asp Leu Lys Pro Thr Asn Ile Leu Leu Gly Asp  
 85 90 95  
 Glu Gly Gln Pro Val Leu Met Asp Leu Gly Ser Met Asn Gln Ala Cys  
 100 105 110  
 Ile His Val Glu Gly Ser Arg Gln Ala Leu Thr Leu Gln Asp Trp Ala  
 115 120 125



Ala Gln Arg Cys Thr Ile Ser Tyr Arg Ala Pro Xaa Leu Phe Ser Val  
 130 135 140

Gln Ser  
 145

<210> 11  
 <211> 505  
 <212> PRT  
 <213> Homo sapiens

<400> 11  
 Met Leu Thr Ser Leu Asn Arg Ser Trp Asn Glu Thr Thr Cys Cys Gly  
 1 5 10 15  
 Arg Ala Ser Phe Leu Glu Leu Cys Thr Gly Gln Ile Gly Arg Thr Pro  
 20 25 30  
 Leu Gly Arg Arg Glu Gly Met Glu Asn Leu Lys His Ile Ile Thr Leu  
 35 40 45  
 Gly Gln Val Ile His Lys Arg Cys Glu Glu Met Lys Tyr Cys Lys Lys  
 50 55 60  
 Gln Cys Arg Arg Leu Gly His Arg Val Leu Gly Leu Ile Lys Pro Leu  
 65 70 75 80  
 Glu Met Leu Gln Asp Gln Gly Lys Arg Ser Val Pro Ser Glu Lys Leu  
 85 90 95  
 Thr Thr Ala Met Asn Arg Phe Lys Ala Ala Leu Glu Glu Ala Asn Gly  
 100 105 110  
 Glu Ile Glu Lys Phe Ser Asn Arg Ser Asn Ile Cys Arg Phe Leu Thr  
 115 120 125  
 Ala Ser Gln Asp Lys Ile Leu Phe Lys Asp Val Asn Arg Lys Leu Ser  
 130 135 140  
 Asp Val Trp Lys Glu Leu Ser Leu Leu Leu Gln Val Glu Gln Arg Met  
 145 150 155 160  
 Pro Val Ser Pro Ile Ser Gln Gly Ala Ser Trp Ala Gln Glu Asp Gln  
 165 170 175  
 Gln Asp Ala Asp Glu Asp Arg Arg Ala Phe Gln Met Leu Arg Arg Asp  
 180 185 190  
 Asn Glu Lys Ile Glu Ala Ser Leu Arg Arg Leu Glu Ile Asn Met Lys  
 195 200 205  
 Glu Ile Lys Glu Thr Leu Arg Gln Tyr Leu Pro Pro Lys Cys Met Gln  
 210 215 220  
 Glu Ile Pro Gln Glu Gln Ile Lys Glu Ile Lys Lys Glu Gln Leu Ser  
 225 230 235 240  
 Gly Ser Pro Trp Ile Leu Leu Arg Glu Asn Glu Val Ser Thr Leu Tyr  
 245 250 255

Lys Gly Glu Tyr His Arg Ala Pro Val Ala Ile Lys Val Phe Lys Lys  
 260 265 270  
 Leu Gln Ala Gly Ser Ile Ala Ile Val Arg Gln Thr Phe Asn Lys Glu  
 275 280 285  
 Ile Lys Thr Met Lys Lys Phe Glu Ser Pro Asn Ile Leu Arg Ile Phe  
 290 295 300  
 Gly Ile Cys Ile Asp Glu Thr Val Thr Pro Pro Gln Phe Ser Ile Val  
 305 310 315 320  
 Met Glu Tyr Cys Glu Leu Gly Thr Leu Arg Glu Leu Leu Asp Arg Glu  
 325 330 335  
 Lys Asp Leu Thr Leu Gly Lys Arg Met Val Leu Val Leu Gly Ala Ala  
 340 345 350  
 Arg Gly Leu Tyr Arg Leu His His Ser Glu Ala Pro Glu Leu His Gly  
 355 360 365  
 Lys Ile Arg Ser Ser Asn Phe Leu Val Thr Gln Gly Tyr Gln Val Lys  
 370 375 380  
 Leu Ala Gly Phe Glu Leu Arg Lys Thr Gln Thr Ser Met Ser Leu Gly  
 385 390 395 400  
 Thr Thr Arg Glu Lys Thr Asp Arg Val Lys Ser Thr Ala Tyr Leu Ser  
 405 410 415  
 Pro Gln Glu Leu Glu Asp Val Phe Tyr Gln Tyr Asp Val Lys Ser Glu  
 420 425 430  
 Ile Tyr Ser Phe Gly Ile Val Leu Trp Glu Ile Ala Thr Gly Asp Ile  
 435 440 445  
 Pro Phe Gln Gly Cys Asn Ser Glu Lys Ile Arg Lys Leu Val Ala Val  
 450 455 460  
 Lys Arg Gln Gln Glu Pro Leu Gly Glu Asp Cys Pro Ser Glu Leu Arg  
 465 470 475 480  
 Glu Ile Ile Asp Glu Cys Arg Ala Ala Gly Arg Leu Val Pro Arg Ser  
 485 490 495  
 Val Ala Ala Ala Arg Ala Val Asp Val  
 500 505

<210> 12  
 <211> 499  
 <212> PRT  
 <213> Homo sapiens

<400> 12  
 Phe Pro Leu Asp Val Glu Tyr Gly Gly Pro Asp Arg Arg Cys Pro Pro  
 1 5 10 15

Pro Pro Tyr Pro Lys His Leu Leu Leu Arg Ser Lys Ser Glu Gln Tyr  
 20 25 30  
 Asp Leu Asp Ser Leu Cys Ala Gly Met Glu Gln Ser Leu Arg Ala Gly  
 35 40 45  
 Pro Asn Glu Pro Glu Gly Gly Asp Lys Ser Arg Lys Ser Ala Lys Gly  
 50 55 60  
 Asp Lys Gly Gly Lys Asp Lys Lys Gln Ile Gln Thr Ser Pro Val Pro  
 65 70 75 80  
 Val Arg Lys Asn Ser Arg Asp Glu Glu Lys Arg Glu Ser Arg Ile Lys  
 85 90 95  
 Ser Tyr Ser Pro Tyr Ala Phe Lys Phe Phe Met Glu Gln His Val Glu  
 100 105 110  
 Asn Val Ile Lys Thr Tyr Gln Gln Lys Val Asn Arg Arg Leu Gln Leu  
 115 120 125  
 Glu Gln Glu Met Ala Lys Ala Gly Leu Cys Glu Ala Glu Gln Glu Gln  
 130 135 140  
 Met Arg Lys Ile Leu Tyr Gln Lys Glu Ser Asn Tyr Asn Arg Leu Lys  
 145 150 155 160  
 Arg Ala Lys Met Asp Lys Ser Met Phe Val Lys Ile Lys Thr Leu Gly  
 165 170 175  
 Ile Gly Ala Phe Gly Glu Val Cys Leu Ala Cys Lys Val Asp Thr His  
 180 185 190  
 Ala Leu Tyr Ala Met Lys Thr Leu Arg Lys Lys Asp Val Leu Asn Arg  
 195 200 205  
 Asn Gln Val Ala His Val Lys Ala Glu Arg Asp Ile Leu Ala Glu Ala  
 210 215 220  
 Asp Asn Glu Trp Val Val Lys Leu Tyr Tyr Ser Phe Gln Asp Lys Asp  
 225 230 235 240  
 Ser Leu Tyr Phe Val Met Asp Tyr Ile Pro Gly Gly Asp Met Met Ser  
 245 250 255  
 Leu Leu Ile Arg Met Glu Val Phe Pro Glu His Leu Ala Arg Phe Tyr  
 260 265 270  
 Ile Ala Glu Leu Thr Leu Ala Ile Glu Ser Val His Lys Met Gly Phe  
 275 280 285  
 Ile His Arg Asp Ile Lys Pro Asp Asn Ile Leu Ile Asp Leu Asp Gly  
 290 295 300  
 His Ile Lys Leu Thr Asp Phe Gly Leu Cys Thr Gly Phe Arg Trp Thr  
 305 310 315 320  
 His Asn Ser Lys Tyr Tyr Gln Lys Gly Ser His Val Arg Gln Asp Ser  
 325 330 335

Met Glu Pro Ser Asp Leu Trp Asp Asp Val Ser Asn Cys Arg Cys Gly  
 340 345 350

Asp Arg Leu Lys Thr Leu Glu Gln Arg Ala Arg Lys Gln His Gln Arg  
 355 360 365

Cys Leu Ala His Ser Leu Val Gly Thr Pro Asn Tyr Ile Ala Pro Glu  
 370 375 380

Val Leu Leu Arg Lys Gly Tyr Thr Gln Leu Cys Asp Trp Trp Ser Val  
 385 390 395 400

Gly Val Ile Leu Phe Glu Met Leu Val Gly Gln Pro Pro Phe Leu Ala  
 405 410 415

Pro Thr Pro Thr Glu Thr Gln Leu Lys Val Ile Asn Trp Glu Asn Thr  
 420 425 430

Leu His Ile Pro Ala Gln Val Lys Leu Ser Pro Glu Ala Arg Asp Leu  
 435 440 445

Ile Thr Lys Leu Cys Cys Ser Ala Asp His Arg Leu Gly Arg Asn Gly  
 450 455 460

Ala Asp Asp Leu Lys Ala His Pro Phe Phe Ser Ala Ile Asp Phe Ser  
 465 470 475 480

Ser Asp Ile Arg Lys His Pro Ala Pro Tyr Val Pro Thr Ile Ser His  
 485 490 495

Pro Met Glu

<210> 13  
 <211> 375  
 <212> DNA  
 <213> Homo sapiens

<400> 13  
 cttgcaggat ttgagttgag gaaaacacag acttccatga gtttgggaac tacgagagaa 60  
 aagacagaca gagtcaaatac tacagcatat ctctcacctc aggaactgga agatgtat 120  
 tatcaatatg atgtaaagtc tgaaatatac agctttggaa tcgtcctctg ggaaatcgcc 180  
 actggagata tcccgtttca aggctgtaat tctgagaaga tccgcaagct ggtggctgtg 240  
 aagcggcagc aggagccact ggggtgaagac tgccttcag agctgcggga gatcattgat 300  
 gagtgccggg cccatgatcc ctctgtgcgg ccctctgtgg atgaaatctt aaagaaactc 360  
 tccacctttt ctaag 375

<210> 14  
 <211> 125  
 <212> PRT  
 <213> Homo sapiens

<400> 14  
 Leu Ala Gly Phe Glu Leu Arg Lys Thr Gln Thr Ser Met Ser Leu Gly  
 1 5 10 15  
 Thr Thr Arg Glu Lys Thr Asp Arg Val Lys Ser Thr Ala Tyr Leu Ser  
 20 25 30

Pro Gln Glu Leu Glu Asp Val Phe Tyr Gln Tyr Asp Val Lys Ser Glu  
35 40 45

Ile Tyr Ser Phe Gly Ile Val Leu Trp Glu Ile Ala Thr Gly Asp Ile  
50 55 60

Pro Phe Gln Gly Cys Asn Ser Glu Lys Ile Arg Lys Leu Val Ala Val  
65 70 75 80

Lys Arg Gln Gln Glu Pro Leu Gly Glu Asp Cys Pro Ser Glu Leu Arg  
85 90 95

Glu Ile Ile Asp Glu Cys Arg Ala His Asp Pro Ser Val Arg Pro Ser  
100 105 110

Val Asp Glu Ile Leu Lys Lys Leu Ser Thr Phe Ser Lys  
115 120 125

<210> 15

<211> 1961

<212> DNA

<213> Homo sapiens

<400> 15

```
tcccgcctgga cgtggagtag ggaggccag accggagggtg cccgcctccg ccctaccga 60
agcacctgct gctgcgcagc aagtcggagc agtacgacct ggacagcctg tgcgcaggca 120
tggagcagag cctccgtgag ggccccaacg agcccgaggc cggcgacaag agccgcaaaa 180
gcgccaaggg ggacaaaggg ggaaggata aaaagcagat tcagacctct cccgttcccg 240
tccgcaaaaa cagcagagac gaagagaaga gagagtcacg catcaagagc tactcgccat 300
acgcctttaa gttcttcag gagcagcacg tggagaatgt catcaaaacc taccagcaga 360
aggttaaccg gaggtgcag ctggagcaag aaatggccaa agctggactc tgtgaagctg 420
agcaggagca gatgcggaag atcctctacc agaaagagtc taattacaac aggttaaga 480
gggccaagat ggacaagtct atgtttgtca agatcaaaac cctggggatc ggtgcctttg 540
gagaagtgtg ccttgcttgt aaggtggaca ctacgcctc gtacgccatg aagaccctaa 600
ggaaaaagga tgtcctgaac cggaatcagg tggcccacgt caaggccgag agggacatcc 660
tggccgaggg agacaatgag tgggtggtca aactctacta ctcttccaa gacaaagaca 720
gcctgtactt tgtgatggac tacatccctg gtggggacat gatgagcctg ctgatccgga 780
tggaggtctt ccctgagcac ctggcccggg tctacatcgc agagctgact ttggccattg 840
agagtgtcca caagatgggc ttcattccacc gagacatcaa gcctgataac attttgatag 900
atctggatgg tcacattaaa ctcacagatt tcggcctctg cactgggttc aggtggactc 960
acaattccaa atattaccag aaaggagacc atgtcagaca ggacagcatg gagccagcg 1020
acctctggga tgatgtgtct aactgtcggg gtggggagac gctgaagacc ctgagcaga 1080
gggcgaggaa gcagcaccag aggtgcctgg cacattcact ggtggggact ccaaaactaca 1140
tcgcacccga ggtgctcctc cgcaagggtt aactcaact ctgtgactgg tggagtgttg 1200
gagtgtattc cttcgagatg ctggtggggc agccgccctt ttgggcacct actcccacag 1260
aaaccagct gaaggtgac aactgggaga acacgctcca cattccagcc caggtgaagc 1320
tgagccctga ggccagggac ctcacacca agctgtgctg ctccgcagac caccgcctgg 1380
ggcggaatgg ggccgatgac ctgaaggccc accccttctt cagcgccatt gacttctcca 1440
gtgacatccg gaagcatcca gcccctacg ttcaccacat cagccacccc atggacacct 1500
cgaatttcga ccccgtagat gaagaaagcc cttggaacga tgccagcgaa ggtagacca 1560
aggcctggga cacactcacc tcgccaata acaagcatcc tgagcagcga ttttacgaat 1620
tcaccttccg aaggttcttt gatgacaatg gctacccctt tcgatgcccc aagccttcag 1680
gagcagaagc ttcacaggct gagagctcag atttagaaaag ctctgatctg gtggatcaga 1740
ctgaaggctg ccagcctgtg tacgtgtaga tgggggccag gcacccccac cactcgtctg 1800
ctcccaggtc aggttcccg agccggtgcc ctcacaggcc aatagggaag ccgagggtg 1860
ttttgttta aattagtccg tcgattactt cacttgaat tctgctcttc accaagaaaa 1920
cccaaacagg acacttttga aaacagcggg gccgcgaatt c 1961
```

<210> 16  
 <211> 588  
 <212> PRT  
 <213> Homo sapiens

<400> 16  
 Pro Leu Asp Val Glu Tyr Gly Gly Pro Asp Arg Arg Cys Pro Pro Pro  
 1 5 10 15  
 Pro Tyr Pro Lys His Leu Leu Leu Arg Ser Lys Ser Glu Gln Tyr Asp  
 20 25 30  
 Leu Asp Ser Leu Cys Ala Gly Met Glu Gln Ser Leu Arg Ala Gly Pro  
 35 40 45  
 Asn Glu Pro Glu Gly Gly Asp Lys Ser Arg Lys Ser Ala Lys Gly Asp  
 50 55 60  
 Lys Gly Gly Lys Asp Lys Lys Gln Ile Gln Thr Ser Pro Val Pro Val  
 65 70 75 80  
 Arg Lys Asn Ser Arg Asp Glu Glu Lys Arg Glu Ser Arg Ile Lys Ser  
 85 90 95  
 Tyr Ser Pro Tyr Ala Phe Lys Phe Phe Met Glu Gln His Val Glu Asn  
 100 105 110  
 Val Ile Lys Thr Tyr Gln Gln Lys Val Asn Arg Arg Leu Gln Leu Glu  
 115 120 125  
 Gln Glu Met Ala Lys Ala Gly Leu Cys Glu Ala Glu Gln Glu Gln Met  
 130 135 140  
 Arg Lys Ile Leu Tyr Gln Lys Glu Ser Asn Tyr Asn Arg Leu Lys Arg  
 145 150 155 160  
 Ala Lys Met Asp Lys Ser Met Phe Val Lys Ile Lys Thr Leu Gly Ile  
 165 170 175  
 Gly Ala Phe Gly Glu Val Cys Leu Ala Cys Lys Val Asp Thr His Ala  
 180 185 190  
 Leu Tyr Ala Met Lys Thr Leu Arg Lys Lys Asp Val Leu Asn Arg Asn  
 195 200 205  
 Gln Val Ala His Val Lys Ala Glu Arg Asp Ile Leu Ala Glu Ala Asp  
 210 215 220  
 Asn Glu Trp Val Val Lys Leu Tyr Tyr Ser Phe Gln Asp Lys Asp Ser  
 225 230 235 240  
 Leu Tyr Phe Val Met Asp Tyr Ile Pro Gly Gly Asp Met Met Ser Leu  
 245 250 255  
 Leu Ile Arg Met Glu Val Phe Pro Glu His Leu Ala Arg Phe Tyr Ile  
 260 265 270  
 Ala Glu Leu Thr Leu Ala Ile Glu Ser Val His Lys Met Gly Phe Ile  
 275 280 285

His Arg Asp Ile Lys Pro Asp Asn Ile Leu Ile Asp Leu Asp Gly His  
 290 295 300  
 Ile Lys Leu Thr Asp Phe Gly Leu Cys Thr Gly Phe Arg Trp Thr His  
 305 310 315 320  
 Asn Ser Lys Tyr Tyr Gln Lys Gly Ser His Val Arg Gln Asp Ser Met  
 325 330 335  
 Glu Pro Ser Asp Leu Trp Asp Asp Val Ser Asn Cys Arg Cys Gly Asp  
 340 345 350  
 Arg Leu Lys Thr Leu Glu Gln Arg Ala Arg Lys Gln His Gln Arg Cys  
 355 360 365  
 Leu Ala His Ser Leu Val Gly Thr Pro Asn Tyr Ile Ala Pro Glu Val  
 370 375 380  
 Leu Leu Arg Lys Gly Tyr Thr Gln Leu Cys Asp Trp Trp Ser Val Gly  
 385 390 395 400  
 Val Ile Leu Phe Glu Met Leu Val Gly Gln Pro Pro Phe Leu Ala Pro  
 405 410 415  
 Thr Pro Thr Glu Thr Gln Leu Lys Val Ile Asn Trp Glu Asn Thr Leu  
 420 425 430  
 His Ile Pro Ala Gln Val Lys Leu Ser Pro Glu Ala Arg Asp Leu Ile  
 435 440 445  
 Thr Lys Leu Cys Cys Ser Ala Asp His Arg Leu Gly Arg Asn Gly Ala  
 450 455 460  
 Asp Asp Leu Lys Ala His Pro Phe Phe Ser Ala Ile Asp Phe Ser Ser  
 465 470 475 480  
 Asp Ile Arg Lys His Pro Ala Pro Tyr Val Pro Thr Ile Ser His Pro  
 485 490 495  
 Met Asp Thr Ser Asn Phe Asp Pro Val Asp Glu Glu Ser Pro Trp Asn  
 500 505 510  
 Asp Ala Ser Glu Gly Ser Thr Lys Ala Trp Asp Thr Leu Thr Ser Pro  
 515 520 525  
 Asn Asn Lys His Pro Glu His Ala Phe Tyr Glu Phe Thr Phe Arg Arg  
 530 535 540  
 Phe Phe Asp Asp Asn Gly Tyr Pro Phe Arg Cys Pro Lys Pro Ser Gly  
 545 550 555 560  
 Ala Glu Ala Ser Gln Ala Glu Ser Ser Asp Leu Glu Ser Ser Asp Leu  
 565 570 575  
 Val Asp Gln Thr Glu Gly Cys Gln Pro Val Tyr Val  
 580 585



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C12N 15/55, C07K 16/40,</b> <b>C12N 9/12, G01N 33/50</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/08180</b> <b>(43) International Publication Date:</b> 17 February 2000 (17.02.2000)
<b>(21) International Application Number:</b> PCT/US99/17630 <b>(22) International Filing Date:</b> 03 August 1999 (03.08.1999) <b>(30) Priority Data:</b> 60/095,270 04 August 1998 (04.08.1998) US 60/099,972 11 September 1998 (11.09.1998) US <b>(60) Parent Application or Grant</b> IMMUNEX CORPORATION [/]; (). VIRCA, G., Duke [/]; (). BIRD, Timothy, A. [/]; (). ANDERSON, Dirk, M. [/]; (). MARKEN, John, S. [/]; (). VIRCA, G., Duke [/]; (). BIRD, Timothy, A. [/]; (). ANDERSON, Dirk, M. [/]; (). MARKEN, John, S. [/]; (). JONES, Simone, L. ; ().		<b>Published</b>
<b>(54) Title: HUMAN cDNAs ENCODING POLYPEPTIDES HAVING KINASE FUNCTIONS</b> <b>(54) Titre: ADNc HUMAINS CODANT DES POLYPEPTIDES A FONCTIONS KINASENS</b>  <b>(57) Abstract</b> <p>The invention is directed to purified and isolated human polypeptides having kinase function, the nucleic acids encoding such polypeptides, processes for production of recombinant forms of such polypeptides, antibodies generated against these polypeptides, fragmented peptides derived from these polypeptides, the use of such polypeptides and fragmented peptides in phosphorylation reactions and as molecular weight markers, the use of such polypeptides and fragmented peptides as controls for peptide fragmentation, the use of such polypeptides in screening assays, and kits comprising these reagents.</p> <b>(57) Abrégé</b> <p>La présente invention concerne des polypeptides humains isolés et purifiés à fonction kinase, les acides nucléiques codant ces polypeptides, des procédés de production de formes recombinées de ces polypeptides, des anticorps produits contre ces polypeptides, et des fragments de peptides obtenus à partir de ces polypeptides. Par ailleurs, cette invention concerne l'utilisation de ces polypeptides et des peptides fragmentés dans des réactions de phosphorylation et comme étalons de masse moléculaire, l'utilisation de ces polypeptides et des peptides fragmentés comme régulateurs de la fragmentation des peptides, l'utilisation de ces polypeptides dans des essais de dépistage, et enfin, des kits renfermant ces réactifs.</p>		



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/55, 9/12, C07K 16/40, G01N 33/50</b>		A3	(11) International Publication Number: <b>WO 00/08180</b>
			(43) International Publication Date: 17 February 2000 (17.02.00)
(21) International Application Number: PCT/US99/17630			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 3 August 1999 (03.08.99)			
(30) Priority Data: 60/095,270 4 August 1998 (04.08.98) US 60/099,972 11 September 1998 (11.09.98) US			
(71) Applicant (for all designated States except US): IMMUNEX CORPORATION [US/US]; 51 University Street, Seattle, WA 98101 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): VIRCA, G., Duke [US/US]; 16690 S.E. 50th Place, Bellevue, WA 98006 (US). BIRD, Timothy, A. [GB/US]; 10804 Olallie Drive N.E., Bainbridge Island, WA 98110 (US). ANDERSON, Dirk, M. [US/US]; 3616 N.W. 64th Street, Seattle, WA 98107 (US). MARKEN, John, S. [US/US]; 11804 9th Avenue S.W., Seattle, WA 98146 (US).			
(74) Agent: JONES, Simone, L.; Immunex Corporation, Law Dept., 51 University Street, Seattle, WA 98101 (US).			Published With international search report.
			(88) Date of publication of the international search report: 29 June 2000 (29.06.00)
(54) Title: HUMAN cDNAs ENCODING POLYPEPTIDES HAVING KINASE FUNCTIONS			
(57) Abstract <p>The invention is directed to purified and isolated human polypeptides having kinase function, the nucleic acids encoding such polypeptides, processes for production of recombinant forms of such polypeptides, antibodies generated against these polypeptides, fragmented peptides derived from these polypeptides, the use of such polypeptides and fragmented peptides in phosphorylation reactions and as molecular weight markers, the use of such polypeptides and fragmented peptides as controls for peptide fragmentation, the use of such polypeptides in screening assays, and kits comprising these reagents.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## INTERNATIONAL SEARCH REPORT

International Application No.

PC., JS 99/17630

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N15/55 C12N9/12 C07K16/40 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C12N C07K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DEAK M. ET AL.: "Mitogen- and stress-activated protein kinase-1 (MSK1) is directly activated by MAPK and SAPK2/p38 and may mediate activation of CREB." EMBO JOURNAL, vol. 17, no. 15, 3 August 1998 (1998-08-03), pages 4426-4441, XP002126284 the whole document, especially Fig. 2 ---	1-32,36
X	MARRA M. ET AL.: "The WashU-HHMI Mouse Project." EMBL DATABASE ENTRY MM1263730; ACCESSION NUMBER AA472165,21 June 1997 (1997-06-21), XP002126285 abstract --- -/--	1-16, 23-31

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

## \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

20 December 1999

Date of mailing of the international search report

26.04.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl  
Fax: (+31-70) 340-3016

Authorized officer

Mandl, B

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT, JS 99/17630

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>PIERRAT B. ET AL.: "RSK-B, a novel ribosomal S6 kinase family member, is a CREB kinase under dominant control of p38(alpha) mitogen-activated protein kinase (p38(alpha)MAPK)." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 45, 6 November 1998 (1998-11-06), pages 29661-29671, XP002126286 the whole document -----</p>	<p>1-16, 23-31</p>

# INTERNATIONAL SEARCH REPORT

Int. application No.  
PCT/US 99/17630

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 33,34,35  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
Claims 1-33,35,36 (all partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 33,34,35

Claims 33,34 and 35 refer to antagonists and agonists of the polypeptides without giving a true technical characterization. Moreover, no such specific compounds are defined in the application. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported (Art. 5 and 6 PCT). No search can be carried out for such purely speculative claims whose wording is, in fact, a mere recitation of the results to be achieved.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

and agonists of said polypeptide; and a method of screening a candidate molecule to identify its ability to mimic the biological activity of said polypeptide.

## 5. Claims: 1-36 (all partially)

A nucleic acid molecule comprising the DNA of SEQ.ID.5 or SEQ.ID.13; a vector containing said nucleic acid molecule; a polypeptide encoded by said nucleic acid molecule (as represented by SEQ.ID.11 or SEQ.ID.14); a nucleic acid molecule hybridizing with said nucleic acid molecule; antibodies specific for said polypeptide; a host cell containing said vector; methods for the production of said polypeptide; a method for screening for agonists or antagonists of said polypeptide; antagonists and agonists of said polypeptide; and a method of screening a candidate molecule to identify its ability to mimic the biological activity of said polypeptide.

## 6. Claims: 1-36 (all partially)

A nucleic acid molecule comprising the DNA of SEQ.ID.6; a vector containing said nucleic acid molecule; a polypeptide encoded by said nucleic acid molecule (as represented by SEQ.ID.12); a nucleic acid molecule hybridizing with said nucleic acid molecule; antibodies specific for said polypeptide; a host cell containing said vector; methods for the production of said polypeptide; a method for screening for agonists or antagonists of said polypeptide; antagonists and agonists of said polypeptide; and a method of screening a candidate molecule to identify its ability to mimic the biological activity of said polypeptide.

## 7. Claims: 1-9,11,13-16,19-36 (all partially)

A nucleic acid molecule comprising the DNA of SEQ.ID.15; a vector containing said nucleic acid molecule; a polypeptide encoded by said nucleic acid molecule (as represented by SEQ.ID.16); a nucleic acid molecule hybridizing with said nucleic acid molecule; antibodies specific for said polypeptide; a host cell containing said vector; methods for the production of said polypeptide; a method for screening for agonists or antagonists of said polypeptide; antagonists and agonists of said polypeptide; and a method of screening a candidate molecule to identify its ability to mimic the biological activity of said polypeptide.